

Defining Rules in Cost Spanning Tree Problems Through the Canonical Form

Gustavo Bergantiños and Juan J. Vidal-Puga

NOTA DI LAVORO 97.2004

JUNE 2004

CTN – Coalition Theory Network

Gustavo Bergantiños, *Research Group in Economic Analysis, Universidade de Vigo*
Juan J. Vidal-Puga, *Departamento de Estadística, Universidade de Vigo*

This paper can be downloaded without charge at:

The Fondazione Eni Enrico Mattei Note di Lavoro Series Index:
<http://www.feem.it/Feem/Pub/Publications/WPapers/default.htm>

Social Science Research Network Electronic Paper Collection:
<http://ssrn.com/abstract=XXXXXX>

The opinions expressed in this paper do not necessarily reflect the position of
Fondazione Eni Enrico Mattei

Defining Rules in Cost Spanning Tree Problems Through the Canonical Form

Summary

We define the canonical form of a cost spanning tree problem. The canonical form has the property that reducing the cost of any arc, the minimal cost of connecting agents to the source is also reduced. We argue that the canonical form is a relevant concept in this kind of problems and study a rule using it. This rule satisfies much more interesting properties than other rules in the literature. Furthermore we provide two characterizations. Finally, we present several approaches to this rule without using the canonical form.

Keywords: Cost spanning tree, Rules, Canonical form

JEL Classification: C71, D70, D85

We thank Hervé Moulin and Stef Tijs for helpful comments. Financial support from the Ministerio de Ciencia y Tecnología and FEDER through grant BEC2002-04102-C02-01 and Xunta de Galicia through grant PGIDIT03PXIC30002PN is gratefully acknowledged.

Address for correspondence:

Juan J. Vidal-Puga
Departamento de Estadística
Universidade de Vigo
A Xunqueira
Pontevedra 36005
Spain
Phone: +34 986802014
E-mail: vidalpuga@uvigo.es

Defining rules in cost spanning tree problems through the canonical form

1 Introduction

Many problems involving network formation have been studied in the operations research and the economic literature. In operations research two issues have been extensively explored: the design of efficient algorithms and the computational complexity. The economic literature focuses on aspects such

like the cost sharing of the network and the design of mechanisms trying to explain the way in which the network forms.

In this paper we focus in the cost sharing aspect. Our contribution can be considered in the well-known literature of cost allocation. We assume that there are no external forces (for example, the market) which determine the final allocation. Agents can achieve agreements directly among themselves, or indirectly by letting the final decision to a neutral referee. In both cases the important issue is to find a "fair allocation" of the cost.

In particular we study cost spanning tree problems (*cstp*). Consider that a group of agents, located at different geographical places, want some particular service which can only be provided by a common supplier, called the source. Agents will be served through connections which entail some cost. However, they do not care whether they are connected directly or indirectly to the source.

There are many economic situations that can be modeled in this way. For instance, several towns may draw power from a common power plant, and hence have to share cost of the distribution network. This example appears in Dutta and Kar (2002). Bergantiños and Lorenzo (2003) study a real situation where villagers should pay the cost of constructing pipes from their respective houses to a water supplier.

The literature on *cstp* starts by defining algorithms for constructing minimal cost spanning trees (*mcst*). We can mention, for instance, the papers of Kruskal (1956) and Prim (1957). But constructing an *mcst* is only a part of the problem. Other important issue is how to allocate the cost associated to the *mcst* among agents.

Bird (1976) studies it using game theory. Bird associates to any *cstp* a cooperative game and proposes a cost allocation rule called the Bird's rule (we call it B). Granot and Huberman (1981, 1984) study the core and the nucleolus of the game. Sharkey (1995) survey most of the literature. Kar (2002) study the Shapley value of the cooperative game defined by Bird, which we call K . Recently, Dutta and Kar (2002) propose a new rule, which we call DK .

Feltkamp, Tijs, and Muto (1994a) introduce a rule for *cstp* called Equal Remaining Obligations. Later, Branzei, Moretti, Norde, and Tijs (2003) call it the P - value, which is the name we use, and give an axiomatic characterization of it.

We associate to each *cstp* a canonical form, which is a *cstp* with the property that by reducing the cost of any arc, the total cost of connecting

agents to the source is also reduced. We argue that the canonical form is a relevant concept for this class of problems and, that it, can be used for defining nice rules. The canonical form "coincides" with the minimal network defined in Bird (1976). In this paper we introduce it in a different way and prove, in propositions 1, 2, and 3, new properties of the canonical form. These new properties will be crucial in our results.

We prove that in canonical problems B and K coincide. Then, for each $cstp$ we define a rule φ as K applied to the canonical form associated.

Surprisingly, φ coincides with P . When we start to study this problem we did not know the unpublished paper of Feltkamp *et al* (1994a). In January 2004 we come across the paper of Branzei *et al* (2003), which was available in the web in December 2003. Through this paper we knew the first one. Then, the originality of our paper is not in the rule we study; it is in the way we introduce it, which has no relation with the way in which is defined P , and the results obtained.

This rule satisfies much more interesting properties than B , K , and DK . We also give two characterizations of φ , which are completely unrelated with the characterization given in Branzei *et al* (2003).

We also present four approaches to this rule without using the canonical form. In the first approach, we prove that P coincides with φ .

The other three approaches are defined in this paper. In the second approach we prove that φ is the Shapley value of a game defined in a similar way to Bird (1976). In the third approach, we give a definition of φ considering only the original problem. In the fourth approach, we obtain that φ can be obtained considering the costs as indivisible goods.

Moreover, Bergantiños and Vidal-Puga (2004) prove that φ can be obtained as the equilibrium payoff of a non-cooperative mechanism.

The paper is organized as follows. In Section 2 we give an overview of the paper. In Section 3 we introduce the $cstp$, along with the rules and properties considered in the paper. In Section 4 we introduce the canonical form and the rule φ . In Section 5 we study the properties satisfied by φ and provide two axiomatic characterizations. The other approaches to φ are presented in Section 6. All the proofs are in the Appendix.


2 An overview of the paper

In this section we present, in an informal way, our most relevant results.

We first discuss, in a very simple example, the allocation proposed by B , K , and DK .

The first non-trivial case in $cstp$ occurs when two agents want to be connected to the source and the optimal choice is that one of the agents connect through the other. The next example is a particular case of this situation.

Example 1. There are two agents. The connection cost between agent 1 and the source is 10, between agent 1 and 2 is 2, and between agent 2 and the source is $10 + x$, where $x \geq 0$. This situation can be represented as follows:



We now discuss how these rules work in this example for each $x \geq 0$. If $x = 0$, the problem is symmetric. The three rules coincide with the symmetric allocation $(6, 6)$.

If $x > 0$, the problem is not symmetric. The unique $mcst$ is $\{(0, 1), (1, 2)\}$. We can proceed in two different ways. First, we ignore x because the arc $(0, 2)$ will not be constructed. Second, since the problem is asymmetric we use the information given by x . Next table shows the proposal given by each of the three rules.

	Agent 1	Agent 2
B	10	2
DK	2	10
K	$6 - \frac{x}{2}$	$6 + \frac{x}{2}$

The three rules use the information given by x . B and DK propose the same allocation independently of x . Moreover, small changes in the cost of arc $(0, 2)$ produce large changes in the proposal. We think that this is unfair. What we are saying is that these rules are not continuous functions on C .

K is a continuous function on C . Take $x = 100$, then $K = (-44, 56)$. This means that agent 2 pays 44 units to agent 1 plus the cost of the network. Again we believe that this allocation is unfair. In this case we claim that the rule must be positive, *i.e.* agents cannot get benefits.

Our conclusion, in Example 1, is that it is better not to take into account the information given by x and always propose $(6, 6)$, even the problem is not symmetric. This example suggests that in some cases it could be better to ignore part of the cost of some arcs if we want to obtain fair rules. In this paper we generalize this idea to the class of all *cstp*.

A *cstp* can be characterized through a matrix C where c_{ij} represents the connection cost between agents i and j . We say that a matrix is canonical if reducing the cost of any arc, the cost of connecting all agents to the source is also reduced. Notice that in Example 1, if $x = 0$ the matrix is canonical but if $x > 0$ the matrix is not canonical.

We define an algorithm for associating to each arbitrary matrix C a unique canonical matrix C^* (Proposition 2). Then, we define the canonical form of a problem as the problem obtained when we change the matrix C by its canonical matrix C^* . In Example 1 the matrix of the problem $x = 0$ is the canonical matrix of each problem with $x > 0$.

Canonical matrices are nice mathematical objects as we prove in propositions 1 and 3. The canonical matrix "coincides" with the minimal network defined in Bird (1976). In this paper we define it in a different way. We also obtain new properties, which will be crucial in most of the proofs of our results.

In Example 1 the three rules coincide in the canonical form ($x = 0$). Even though this result is not true in general; we prove that B and K coincide in canonical matrices and DK is different. Then, we define a rule φ for a general matrix C as the rule K (or B) applied to its canonical matrix C^* .

Our next step is to explain why φ is a suitable rule for these problems. We first study which properties are satisfied by φ and compare it with the other three rules.

In Example 1 we criticize the proposals of B , DK , and K in terms of the properties of continuity (*CON*) and positivity (*POS*). Nevertheless, these rules satisfy other interesting properties. B satisfies core selection (*CS*). However B fails to satisfy cost monotonicity (*CM*), which says that if the connection cost between agents i and j increases, none of these agents should

pay less. K satisfies CM but fails to satisfy CS and DK satisfies both.

We also consider other properties as strong cost monotonicity (SCM), population monotonicity (PM), equal sharing of extra cost ($ESEC$), separability (SEP), equal contributions (EC), and independence of irrelevant trees (IIT).

SCM means that if the connection cost between agents i and j increases, no agent should pay less. PM says that if we add new agents to the problem no agent is worse.

Consider a problem where, for any agent, his most expensive cost is to the source. Moreover, the connection cost to the source is the same for each agent. Assume that this connection cost increases x . $ESEC$ says that if agent i pays f_i in the original problem and n is the number of agents, he must pay $f_i + \frac{x}{n}$ when the cost increases.

Two subsets of agents, S and $N \setminus S$, can connect to the source separately or can connect jointly. If there are no savings when they connect jointly, SEP says that agents must pay the same in both circumstances.

EC says that the impact of the connection of agent j on agent's i cost coincides with the impact of the connection of agent i on agent's j cost. IIT says that if two $cstp$ have a common $mcst$, both problems must have the same solution.

The next table summarizes the properties satisfied by the four rules.

	B	K	DK	φ
CON	no	YES	no	YES
POS	YES	no	YES	YES
CS	YES	no	YES	YES
CM	no	YES	YES	YES
SCM	no	no	no	YES
PM	no	no	no	YES
$ESEC$	YES	YES	no	YES
SEP	YES	no	YES	YES
EC	no	no	no	YES
IIT	no	no	no	YES

These results suggest two comments. Firstly, φ satisfies much more interesting properties than the other three rules.

Secondly, φ and K coincide in canonical matrices. φ does not use the information of C which is not in its canonical form and then $\varphi(C) = K(C^*)$.

K uses this information and thus $K(C)$ can be different to $K(C^*)$. If we consider the properties satisfied by both rules, it seems better not to use this information. A similar comment can be applied to φ and B .

Even though we are considering two particular rules, the above suggests that we may define rules in the general problem through its canonical form. A natural question is whether this is a good strategy.

In Proposition 5 (a) we characterize the rules defined through its canonical form as the only rules satisfying *IIT*. In Proposition 5 (b) we prove that if a rule is not defined through the canonical form then it does not satisfy *SCM*. This means that if we do not use the canonical form we miss at least an important property. Is it possible to gain anything? If we focus on B , DK , K , and φ and the properties mentioned here the answer is no. In general, we do not know. Although we believe that this is an interesting question, it is out of the scope of the paper. Here, we only want to point out that the canonical form is an important concept, which should be studied carefully because it can be very influential in the problem.

Later, we present two characterizations of φ . The first one is with *EC* and the second one is with *SCM*, *PM*, and *ESEC*.

Let us comment the second characterization. *SCM* and *PM* are quite standard properties, which are used often in many economic problems. We believe that these properties are very natural and any fair rule should satisfy both. *ESEC* is a property defined explicitly for *cstp*. We believe that is a natural property with a clear meaning but we do not feel that every fair rule should satisfy it. Thus, we can see *ESEC* as a property selecting a rule among the set of "fair rules" (rules satisfying *SCM* and *PM*).

Moreover, the definition of these three properties is unrelated to the canonical form. We believe that this is other argument supporting the canonical form as a good way for defining rules.

Finally, we argue that φ is a "focal point" of this class of problems because it can be achieved from different approaches. We mention four. In three of these approaches we do not use canonical matrices. In the fourth we use it but in a different way.

The first approach is through the *P - value*. As we said before the *P - value* was introduced in Feltkamp *et al* (1994a) in a very different way. In this paper we prove that φ coincides with *P*.

Second approach. Bird (1976) associated to each problem C a characteristic function v where $v(S)$ denotes the cost of connecting agents in S to the source assuming that agents of $N \setminus S$ are not present. Since the Shapley value is a suitable concept for TU games we can use it in *cstp*. Kar (2002) gives additional arguments supporting the Shapley value of v . He characterizes it considering only the problem C .

Following Bird's approach we associate to each problem C a characteristic function v^+ where $v^+(S)$ denotes the cost of connecting agents in S to the source *assuming that agents of $N \setminus S$ are already connected*. Surprisingly, the Shapley value of v^+ coincides with φ .

The third approach is looking only to the problem C . We explain it with an example. Assume that $\{(0, 1), (1, 2), (2, 3)\}$ is the unique *mcst*. Moreover, the cost of its arcs are 12, 4, and 6, respectively. We address the problem of how to divide the cost of each arc between the agents. We define a rule through an algorithm in which, at each step, we select for each agent both an arc and the part of the cost of this arc he must pay for.

Step 1. Every agent selects the cheapest arc to which he belongs. Thus agents 1 and 2 select $(1, 2)$ and agent 3 selects $(2, 3)$. Since agents 1 and 2 select arc $(1, 2)$ this cost is equally divided between them. Agent 3 pays half of the cost of arc $(2, 3)$.

Step 2. Agents 1 and 2 select the cheapest non-paid arc to which one of them belong. Thus, both select $(2, 3)$. Agent 3 selects the cheapest non-paid arc to which he belongs, *i.e.* $(2, 3)$. Then, the non-paid cost of arc $(2, 3)$ is equally divided among the three agents.

Step 3. The three agents select arc $(0, 1)$, the only whose cost was not paid. This cost is divided equally among agents.

Then, agents 1 and 2 pay $\frac{3}{6}4 + \frac{1}{6}6 + \frac{2}{6}12 = 7$ and agent 3 pays $\frac{3}{6}6 + \frac{1}{6}6 + \frac{2}{6}12 = 8$.

Fourth approach. In B and DK we first select a tree minimizing the cost of connecting agents to the network. Then, we assign the whole cost of each arc to one agent following some specific protocol. Thus, we can consider B and DK as rules assigning indivisible goods (cost of the arcs).

In the case of B we can interpret it assuming that agents sequentially connect to one of the cheapest available options. Each agent pays his connection cost. This protocol can provoke unfair allocations as we can see in Example 1 when x is very small.

A classical way for recovering fairness from an order-depending allocation is to take the average over the set of all permutations. In general problems,

this approach is incompatible with efficiency. Nevertheless, if C is a general problem and C^* is its canonical matrix, it is possible to generate an efficient and fair allocation for C averaging over the permutations in C^* . This is the fourth approach.

3 The cost spanning tree problem

This section is divided in three subsections. In the first subsection, we introduce the problem. In the second subsection, we introduce some rules of the literature. Finally, in the third subsection, we present some properties of the rules.

3.1 The problem

Let $\mathcal{N} = \{1, 2, \dots\}$ be the set of all possible agents. We are interested in networks whose nodes are elements of a set $N_0 = N \cup 0$, where $N \subset \mathcal{N}$ is finite and 0 is a special node called the *source*. Usually we take $N = \{1, \dots, n\}$. Our interest lies on networks where each node in N is (directly or indirectly) connected to the source.

Let Π_N be the set of all permutations over the finite set N . Given $\pi \in \Pi_N$, let $Pre(i, \pi)$ denote the set of elements of N which come before i in the order given by π , *i.e.* $Pre(i, \pi) = \{j \in N \mid \pi(j) < \pi(i)\}$. Given $S \subset N$, let π_S denote the order induced by π among agents in S .

A *cost matrix* $C = (c_{ij})_{i,j \in N_0}$ on N represents the cost of direct link between any pair of nodes. We assume that $c_{ij} = c_{ji} \geq 0$ for each $i, j \in N_0$ and $c_{ii} = 0$ for each $i \in N_0$. Since $c_{ij} = c_{ji}$ we will work with undirected arcs, *i.e.* $(i, j) = (j, i)$.

We denote by \mathcal{C}^N the set of all cost matrices on N . Given $C, C' \in \mathcal{C}^N$ we say $C \leq C'$ if $c_{ij} \leq c'_{ij}$ for all $i, j \in N_0$.

A *cost spanning tree problem*, briefly *cstp*, is a pair (N_0, C) where $N \subset \mathcal{N}$ is the set of agents, 0 is the source, and $C \in \mathcal{C}^N$ is the cost matrix.

A *network* g over N_0 is a subset of $\{(i, j) \text{ such that } i, j \in N_0\}$. The elements of g are called *arcs*.

Given a network g and a pair of nodes i and j , a *path* from i to j in g is a sequence of arcs $\{(i_{h-1}, i_h)\}_{h=1}^l$ satisfying $(i_{h-1}, i_h) \in g$ for all $h \in \{1, 2, \dots, l\}$, $i = i_0$ and $j = i_l$. We say that $i, j \in N$ are *linked* in g if there exists a path

from i to j which does not include the source. If $(i, j) \in g$, we say that i and j are *directly linked* in g . We say that the node i is *connected* in the network g if there exists a path from i to the source. Otherwise we say that i is *unconnected* in g .

Given a network g and $S \subset N$ we denote by g_S the network induced by g among agents in S , *i.e.*

$$g_S = \{(i, j) \in g \text{ such that } \{i, j\} \subset S\}.$$

A *tree* is a network satisfying that for all $i \in N$ there is a unique path from i to the source. If t is a tree we usually write $t = \{(j, j^0)\}_{j \in N}$ where j^0 represents the first agent in the unique path in t from j to 0.

We denote by \mathcal{G}^N the set of all networks over N_0 and by \mathcal{G}_0^N the set of networks such that every agent in N is connected to the source.

Given a *cstp* (N_0, C) and $g \in \mathcal{G}^N$, we define the *cost* associated to g as

$$c(N_0, C, g) = \sum_{(i,j) \in g} c_{ij}.$$

When there are no ambiguities, we write $c(g)$ or $c(C, g)$ instead of $c(N_0, C, g)$.

A *minimum cost spanning tree* for (N_0, C) , briefly a *mcst*, is a tree $t \in \mathcal{G}_0^N$ such that $c(t) = \min_{g \in \mathcal{G}_0^N} c(g)$. It is well-known in the literature about *cstp* that

there exists a *mcst*, even though it does not need to be unique. Given a *cstp* (N_0, C) we denote by $m(N_0, C)$ the cost associated to any *mcst* t in (N_0, C) .

Given a *cstp* (N_0, C) , we define the *cstp* induced by C in $S \subset N$ as (S_0, C) .

Bird (1976) associated to each *cstp* (N_0, C) a *TU* game (N, v_C) where for each coalition $S \subset N$,

$$v_C(S) = m(S_0, C).$$

Usually, we write v instead of v_C .

We now introduce some well-known results of *TU* games, which will be used through the paper. We introduce them considering the *TU* game as a cost game.

We define the *core* of the *TU* game (N, w) as

$$\text{core}(N, w) = \left\{ x \in R^N \mid \sum_{i \in N} x_i = w(N) \text{ and } \forall S \subset N, \sum_{i \in S} x_i \leq w(S) \right\}.$$

We say that (N, w) is *concave* if, for all $S, T \subset N$ and $i \in N$ such that $S \subset T$ and $i \notin T$,

$$w(S \cup i) - w(S) \geq w(T \cup i) - w(T).$$

We denote by $Sh(N, w)$ the *Shapley value* (Shapley, 1953) of the *TU* game (N, w) . It is well-known that the Shapley value belongs to the core when the game is concave.

3.2 Rules

One of the most important issues addressed in the literature about *cstp* is how to divide the cost of connecting agents to the source among them. We now introduce, briefly, some of the rules studied in the literature.

A (*cost allocation*) *rule* is a function ψ such that $\psi(N_0, C) \in \mathbb{R}^N$ for each *cstp* (N_0, C) and $\sum_{i \in N} \psi_i(N_0, C) = m(N_0, C)$. As usually, $\psi_i(N_0, C)$ represents the cost assigned to agent i .

Notice that we implicitly assume that agents built a *mcst*. As far as we know, all the rules proposed in the literature make this assumption.

Given a *cstp*, Prim (1957) provides an algorithm for solving the problem of connecting all agents to the source such that the total cost of creating the network is minimal. The idea of this algorithm is quite simple: starting from the source we construct a network by consecutively adding arcs with the lowest cost, without introducing cycles.

Formally, Prim's algorithm is defined as follows. We start with $S^0 = \{0\}$ and $g^0 = \emptyset$.

Stage 1: Take the arc $(0, i)$ such that $c_{0i} = \min_{i \in N} \{c_{0i}\}$. If there are several arcs $(0, i)$ satisfying this condition, select one of them. Now, $S^1 = \{0, i\}$ and $g^1 = \{(0, i)\}$.

Assume that we have defined S^p and g^p . We now define S^{p+1} and g^{p+1} . Take an arc (j, i) such that $c_{ji} = \min_{k \in S^p, l \in N_0 \setminus S^p} \{c_{kl}\}$. If there are several arcs (j, i) satisfying this condition, select one of them. Now, $S^{p+1} = S^p \cup \{i\}$ and $g^{p+1} = g^p \cup \{(j, i)\}$.

This process finishes in n stages. We say that g^n is a tree obtained via Prim's algorithm. Notice that this algorithm leads to a tree, but not necessarily unique.

We now introduce four rules of the literature: Bird's rule, P 's rule, Kar's rule and Dutta-Kar's rule.

Bird's rule (Bird, 1976) and Dutta-Kar's rule (Dutta and Kar, 2002) are defined through Prim's algorithm. We first assume that there is a unique tree t obtained through Prim's algorithm.

Let i^0 be the first node in the unique path in t from i to the source. Then, given a $cstp (N_0, C)$, *Bird's rule* (B) is defined for each $i \in N$ as

$$B_i(N_0, C) = c_{i^0i}.$$

The idea of this rule is quite simple. Agents connect sequentially to the source following Prim's algorithm and each agent pays the cost of his connection.

Dutta-Kar's rule is defined in a more elaborate way. Assume that agents, according with Prim's algorithm, connect in the order $1, 2, \dots, n$. First agent 1 connects to the source. We define $p^1 = c_{01}$. Now agent 2 connects to 2^0 where $c_{2^02} = \min \{c_{02}, c_{12}\}$. We take $x_1 = \min \{p^1, c_{2^02}\}$ and $p^2 = \max \{p^1, c_{2^02}\}$. Now agent 3 connects to 3^* where $c_{3^03} = \min \{c_{03}, c_{13}, c_{23}\}$. We take $x_2 = \min \{p^2, c_{3^03}\}$ and $p^3 = \max \{p^2, c_{3^03}\}$. This process continue until agent n . In this case we take $x_n = \max \{p^{n-1}, c_{n^0n}\}$. See Dutta and Kar (2002) for a formal definition.

Dutta-Kar's rule (DK) is defined for each $i \in N$ as

$$DK_i(N_0, C) = x_i.$$

Assume now that the tree associated to Prim's algorithm is not unique. In this case Bird's rule and Dutta-Kar's rule can be defined as an average over the trees associated to Prim's algorithm.

Dutta and Kar (2002) proceed as follows. Given $\pi \in \Pi_N$ they define $DK^\pi(N_0, C)$ as the allocation obtained when we apply the previous protocol to (N_0, C) and solving the indifferences selecting the first agent given by π . Then, they define

$$DK(N_0, C) = \frac{1}{n!} \sum_{\pi \in \Pi_N} DK^\pi(N_0, C).$$

$B(N_0, C)$ can be defined in a similar way.

Kruskal (1956) introduced an algorithm for computing the *mest* of a *cstp*. Feltkamp *et al* (1994a) define a rule through Kruskal’s algorithm. This rule was called Equal Remaining Obligations (*ERO*) in this paper. Recently, Branzei *et al* (2003) call it the *P – value* (*P*). We give the idea of this rule following Branzei *et al* (2003). Initially, every agent has an obligation 1 and the network is empty. We now apply Kruskal’s algorithm and the obligations of the agents decrease when we add an arc to the network. This obligation is $\frac{1}{n_i}$, where n_i is the number of agents connected directly or indirectly to agent i through the network. At each step of the algorithm each agent pays the part of the cost of the arc we add given by the difference between his obligation before adding the arc and his obligation after adding the arc. See Branzei *et al* (2003) for a formal definition.

Another approach for defining rules is using game theory. Bird (1976) associated a cooperative game with any *cstp*. Later, several authors defined rules to the *cstp* using this cooperative game. For instance, Granot and Huberman (1981, 1984) studied the core and the nucleolus, and Kar (2002) studied the Shapley value.

Kar’s rule (*K*) is defined as

$$K(N_0, C) = Sh(N, v).$$

3.3 Properties

We now introduce some properties of rules.

We say that a rule ψ satisfies:

Core selection (*CS*) if for all *cstp* (N_0, C) and all $S \subset N$,

$$\sum_{i \in S} \psi_i(N_0, C) \leq m(S_0, C).$$

This property says that no group of agents can be better constructing their own network instead of paying what the rule ψ proposes to them. Notice that core selection is equivalent to say that $\psi(N_0, C) \in core(N, v_C)$.

Cost monotonicity (CM) if for all *cstp* (N_0, C) and (N_0, C') such that $c_{ij} < c'_{ij}$ for some $i, j \in N_0$ and $c'_{kl} = c_{kl}$ otherwise,

$$\psi_i(N_0, C) \leq \psi_i(N_0, C').$$

Cost monotonicity implies that if the connection cost between agents i and j increases and the rest of costs are the same, then agents i and j cannot be better.

Strong cost monotonicity (SCM) if for all *cstp* (N_0, C) and (N_0, C') such that $c_{ij} < c'_{ij}$ for some $i, j \in N_0$ and $c'_{kl} = c_{kl}$ otherwise,

$$\psi_k(N_0, C) \leq \psi_k(N_0, C') \text{ for all } k \in N.$$

Strong cost monotonicity implies that if the connection cost between agents i and j increases and the rest of the costs remain the same, then no agent can be better.

Population monotonicity (PM) if for all *cstp* (N_0, C) , all $S \subset N$, and all $i \in S$,

$$\psi_i(N_0, C) \leq \psi_i(S_0, C).$$

Population monotonicity says that if new agents join a society no agent of the initial society can be worse.

Continuity (CON) if ψ is a continuous function of C . This property says that small changes in the connection costs of agents cannot provoke large changes in the amount they have to pay.

Positivity (POS) if for all *cstp* (N_0, C) and all $i \in N$

$$\psi_i(N_0, C) \geq 0.$$

This property says that agents cannot get benefits.

Separability (SEP) if for all *cstp* (N_0, C) and $S \subset N$ satisfying $m(N_0, C) = m(S_0, C) + m((N \setminus S)_0, C)$,

$$\psi_i(N_0, C) = \begin{cases} \psi_i(S_0, C) & \text{if } i \in S \\ \psi_i((N \setminus S)_0, C) & \text{if } i \in N \setminus S. \end{cases}$$

Two subset of agents, S and $N \setminus S$, can connect to the source separately or can connect jointly. If there is not savings when they connect jointly, separability says that agents must pay the same in both circumstances.

This property appears in Megiddo (1978), Granot and Huberman (1981), and Granot and Maschler (1998). They use the name decomposition, instead of separability, and study it relation with the core and the nucleolus of (N, v_C) . For instance, Granot and Huberman (1981) prove that the core and the nucleolus satisfy *SEP*.

Equal Sharing of Extra Cost (ESEC). Let (N_0, C) and (N_0, C') be such that $c_{0i} = c_0$ and $c'_{0i} = c'_0$ for all $i \in N$, $c_0 < c'_0$, and $c_{ij} = c'_{ij} \leq c_0$ for all $\{i, j\} \subset N$. Then,

$$\psi_i(N_0, C') = \psi_i(N_0, C) + \frac{c'_0 - c_0}{n}.$$

A group of agents face a problem (N_0, C) in which all of them have the same connection cost to the source ($c_{i0} = c_0$) and this cost is larger than the connection costs between agents ($c_{ij} \leq c_0$). Under these circumstances, in all *mcst* one agent (any of them) connects to the source directly, and the rest connect to the source through this agent. Moreover, they agree that the right solution is $\psi(N_0, C)$. Assume that there was a mistake and the connection cost to the source is $c'_0 > c_0$. *ESEC* says that agents must pay equally this extra cost $c'_0 - c_0$.

The next properties are more technical but still very reasonable.

We say that two *cstp* (N_0, C) and (N_0, C') are *directly equivalent* if there exists a tree t such that t is a *mcst* for both (N_0, C) and (N_0, C') and it satisfies $c_{ij} = c'_{ij}$ for all $(i, j) \in t$.

We say that (N_0, C) and (N_0, C') are *equivalent* if there exists a succession of cost matrices $\{C^0, C^1, \dots, C^p\}$ with $C^0 = C$, $C^p = C'$, and (N_0, C^{q-1}) directly equivalent to (N_0, C^q) for each $q = 1, 2, \dots, p$.

Notice that if (N_0, C) and (N_0, C') are equivalent then $m(N_0, C) = m(N_0, C')$.

A rule ψ satisfies *Independence of Irrelevant Trees (IIT)* if $\psi(N_0, C) = \psi(N_0, C')$ whenever (N_0, C) and (N_0, C') are equivalent.

Remark 1. Dutta and Kar (2002) define the property of *tree invariance*. This property says that the rule must depends only on the set of *mcst*. They prove that *DK* satisfies tree invariance.

Notice that if a rule satisfies *IIT* it also satisfies tree invariance. However, tree invariance does not imply *IIT*. It is easy to check it in Example 1 taking $x = 0$ and $x = 100$.

Given a *cstp* (N_0, C) and $i \in N$ we denote by $((N \setminus \{i\})_0, C^{+i})$ the *cstp* obtained from (N_0, C) when agents of $N \setminus \{i\}$ have to be connected, assuming that agent i is already connected. This means that for all $j \in N \setminus \{i\}$, $c_{j0}^{+i} = \min \{c_{ij}, c_{j0}\}$ and $c_{jk}^{+i} = c_{jk}$ when $k \neq 0$.

We say that ψ satisfies *Equal Contributions* (*EC*) if for all $i, j \in N, i \neq j$,

$$\psi_i(N_0, C) - \psi_i((N \setminus \{j\})_0, C^{+j}) = \psi_j(N_0, C) - \psi_j((N \setminus \{i\})_0, C^{+i}).$$

EC says that the impact of the connection of agent j on agent's i cost coincides with the impact of the connection of agent i on agent's j cost.

There are some relations among these properties. It is trivial that *SCM* implies *CM* whereas the reciprocal is false.

PM implies *SEP*. Let ψ be a rule satisfying *PM* and $S \subset N$ as in the definition of *SEP*. By *PM* we know that $\psi_i(S_0, C) \geq \psi_i(N_0, C)$ for all $i \in S$ and $\psi_i((N \setminus S)_0, C) \geq \psi_i(N_0, C)$ for all $i \in N \setminus S$. Since $m(N_0, C) = m(S_0, C) + m((N \setminus S)_0, C)$ it is easy to conclude that ψ satisfies *SEP*. The reciprocal is false.

PM implies *CS*. For all $S \subset N$, $\sum_{i \in S} \psi_i(N_0, C) \leq \sum_{i \in S} \psi_i(S_0, C) = m(S_0, C)$.

The reciprocal is false.

In Section 4 (Proposition 5) we prove that *SCM* implies *IIT*.

4 The canonical form of a cost spanning tree problem

In this section we associate to each *cstp* a unique canonical matrix, which has the property that, if we reduce the cost of any arc, then the cost of connecting agents to the source is also reduced. The canonical matrix coincides with the minimal network defined in Bird (1976). We define the canonical form of a *cstp* as the problem obtained by replacing the initial cost matrix by its canonical matrix associated.

We prove in propositions 1 and 3 that canonical matrices have nice mathematical properties, which will be often used in the paper.

Moreover, Bird's rule and Kar's rule coincide in canonical matrices. This allows us to define the rule φ for general *cstp* as Kar's rule (or Bird's rule) of the canonical form of the original problem.

We also study the rules that only depend on the canonical form and we prove that they coincide with the rules satisfying *IIT*. Finally, we obtain that if a rule does not only depend on the canonical form, it does not satisfy strong cost monotonicity. This allows to argue that defining rules through the canonical form is a good approach to *cstp*.

Given a *cstp* (N_0, C) , we say that C is *canonical* if we cannot decrease the cost of an arc without decreasing $m(N_0, C)$.

Remark 2. If C is canonical, for any $i, j \in N_0$ there exists a *mcst* t of (N_0, C) such that $(i, j) \in t$.

In the next proposition we characterize the canonical matrices.

Proposition 1. C is a canonical matrix if and only if there exists a *mcst* t in (N_0, C) satisfying the two following conditions:

- (A1) $t = \{(i_{p-1}, i_p)\}_{p=1}^n$ where $i_0 = 0$ (the source).
- (A2) Given $i_p, i_q \in N_0$, $p < q$ then, $c_{i_p i_q} = \max_{p < r \leq q} \{c_{i_{r-1} i_r}\}$.

Proof. See the appendix.

This means that canonical matrices are uniquely determined by a *mcst* t .

Given a *cstp* (N_0, C) we say that C^* is a *canonical matrix associated to* C if C^* is a canonical matrix, $C \geq C^*$, and $m(N_0, C) = m(N_0, C^*)$. Later we will prove that any *cstp* has a unique associated canonical matrix.

We now introduce an algorithm for associating to each arbitrary matrix $C \in \mathcal{C}^N$ a canonical matrix C^* . This algorithm has $n + 1$ stages and it is inspired by Prim's algorithm.

Let $t^0 = \{(i^0, i)\}_{i \in N}$ be a *mcst* in (N_0, C) . Remember that i^0 is the first agent in the unique path in t from i to 0. Initially, take $C^0 = C$, $S^0 = \{0\}$, and $g^0 = \emptyset$.

Stage 1. Define $T_1 = \{i \in N \mid i^0 = 0\}$. Since t^0 is a tree, $T_1 \neq \emptyset$. Take an arc $(0, i_1)$ such that $i_1 \in T_1$ and $c_{0i_1} = \min_{i \in T_1} \{c_{0i}\}$. If there are several arcs satisfying this condition, take one of them. Now $C^1 = C^0$, $S^1 = \{0, i_1\}$, and $g^1 = \{(0, i_1)\}$.


Assume that we have defined Stage r for all $r \leq p - 1$. We now define Stage p when $p \leq n$.

Stage p . We define $T_p = \{i \in N \mid i \in N_0 \setminus S^{p-1} \text{ and } i^0 \in S^{p-1}\}$. Since t^0 is a tree, $T_p \neq \emptyset$. Take an arc (i_p^0, i_p) such that $c_{i_p^0 i_p} = \min_{i \in T_p} \{c_{i^0 i}\}$. If there are several arcs satisfying this condition, take one of them. We define C^p such that $c_{kj}^p = c_{kj}^{p-1}$ if $(k, j) \neq (i_{p-1}, i_p)$ and $c_{i_{p-1} i_p}^p = c_{i_{p-1} i_p}^0$. Moreover, $S^p = S^{p-1} \cup \{i_p\}$ and $g^p = g^{p-1} \cup \{(i_{p-1}, i_p)\}$.

Stage $n + 1$. We define C^* such that $c_{i_p i_q}^* = \max_{p < r \leq q} \{c_{i_{r-1} i_r}^n\}$ for all $i_p, i_q \in N_0$, $p < q$. Moreover $t = g^n$.


If we apply this algorithm to Example 1 we obtain that for all $cstp$ where $x > 0$, the canonical matrix associated corresponds to the $cstp$ where $x = 0$. We now apply this algorithm in a more complicate example.

Example 2. Consider the following $cstp$:




It is trivial to see that there is a unique $mcst$ $\{(0, 1), (1, 2), (1, 3)\}$.

1. Stage 1. We select the arc $(0, 1)$. Then, $g^1 = \{(0, 1)\}$, $S^1 = \{0, 1\}$, and $C^1 = C$.
2. Stage 2. We select the arc $(1, 2)$. Then, $g^2 = \{(0, 1), (1, 2)\}$, $S^2 = \{0, 1, 2\}$, and $C^2 = C^1$.
3. Stage 3. We select the arc $(1, 3)$. Then, $g^3 = \{(0, 1), (1, 2), (2, 3)\}$, $S^3 = \{0, 1, 2, 3\}$, and C^3 is given by



4. Stage 4. $t = \{(0, 1), (1, 2), (2, 3)\}$. Moreover, C^* is given by



Proposition 2. C^* is a canonical matrix, $C \geq C^*$, C^* is equivalent to C , and C^* is the unique canonical matrix associated to C .

Proof: See the Appendix.

As a consequence of this proposition, for any $cstp (N_0, C)$ we can define the canonical form (N_0, C^*) as the $cstp$ where C^* is the unique canonical matrix associated to C .

If we compute the B , DK , and K in Example 2 we obtain

	Agent 1	Agent 2	Agent 3
Bird	12	4	6
Dutta-Kar	4	6	12
Kar	3.5	7.5	11

If we compute B , DK , and K in the canonical form associated to Example 2 we obtain

	Agent 1	Agent 2	Agent 3
Bird	7	7	8
Dutta-Kar	6	6	10
Kar	7	7	8

Notice B and K coincide in the canonical form associated to Example 1. Later we will prove that this is a general result.

In the next proposition we give a list of properties of canonical matrices. These properties will be used often in the rest of the paper. We will assume $t = \{(i-1, i)\}_{i=1}^n$ only to simplify the notation.

Proposition 3. Let (N_0, C) be a *cstp* where C is canonical and $t = \{(i-1, i)\}_{i=1}^n$ is a *mcst* in (N_0, C) satisfying (A1) and (A2). Take $S \subset N$. We can assume that $S = \{i_1, \dots, i_{|S|}\}$, $i_0 = 0$, and $i_{p-1} \leq i_p$ for all $p = 1, \dots, |S|$. Then,

- (a) $t' = \{(i_{p-1}, i_p)\}_{p=1}^{|S|}$ is a *mcst* of (S_0, C) and $v(S) = \sum_{p=1}^{|S|} c_{i_{p-1}i_p}$.
- (b) $v(S) - v(S \setminus \{i_q\}) = \min \{c_{i_{q-1}i_q}, c_{i_q i_{q+1}}\}$ if $i_q \neq i_{|S|}$ and $v(S) - v(S \setminus \{i_{|S|}\}) = c_{i_{|S|-1}i_{|S|}}$.
- (c) (N, v) is concave.

Proof. See the Appendix.

Remark 3. Given a tree t and $(i, j) \in t$, c_{ij} denote the cost of the arc (i, j) . Bird (1976) define the *minimal network* (N_0, C^t) associated to t in the following way. For each $i, j \in N_0$ such that $(i, j) \notin t$ let g_{ij} denote the unique path in t connecting i and j . Then, $c_{ij}^t = c_{ij}$ if $(i, j) \in t$ and $c_{ij}^t = \max_{(k,l) \in g_{ij}} \{c_{kl}\}$ if $(i, j) \notin t$. Bird (1976) uses this minimal network in order to define the irreducible core of a *cstp*, which is a subset of the core.

Let (N_0, C) be a *cstp* and t a *mcst* in (N_0, C) . It is not difficult to prove that C^t coincides with C^* . Then, for all *mcst* t the minimal network associated to t coincides with the canonical form.

Proposition 4. Let (N_0, C) be a *cstp* where C is canonical. Then, $K(N_0, C) = B(N_0, C)$.

Proof. See the Appendix.

Proposition 4 allows the following definition.

Definition 1. Given a *cstp* (N_0, C) we define the rule φ as

$$\varphi(N_0, C) = K(N_0, C^*) = B(N_0, C^*)$$

where C^* is the canonical matrix associated to C .

As we said in the initial discussion of Example 1 we define the rule through the canonical form. Moreover, we define it as K and B because the fact that these rules, which are very different in general, coincide in canonical matrices could mean something. In the next section we will give more arguments justifying this approach.

But now we discuss an important issue of the paper, which is defining rules through the canonical form. We first characterize the rules which depends only on the canonical form, *i.e.* $\psi(N_0, C) = \psi(N_0, C^*)$ for all *cstp* (N_0, C) .

Proposition 5. (a) A rule ψ depends only on the canonical form if and only if ψ satisfies *IIT*.

(b) *SCM* implies *IIT*.

Proof. See the Appendix.

The reciprocal of part (b) is false because there exist rules satisfying *IIT* but not *SCM*. For instance,

$$\psi_i(N_0, C) = \frac{c_{0i}^*}{\sum_{j=1}^n c_{0j}^*} m(N_0, C) \text{ for all } i \in N$$

satisfies *IIT* but not *SCM*.

One of the main messages of the paper is that defining rules through the canonical form is a very good approach for obtaining nice rules in *cstp*. Proposition 5 (b) is a strong argument supporting this approach. If we consider a rule that does not depend on the canonical form, this rule will not satisfy *SCM*. Something will be missing for sure.

An interesting question from our point of view is whether it is possible to define a rule which does not depend on the canonical form, it satisfies

interesting properties, but no rule depending only on the canonical form satisfies these properties. In the next section we will see that this is not a trivial question because this is not the case of the rules studied in the literature.

5 Properties of the rule

In this section we prove that the rule φ satisfies all the properties stated in the paper. If we compare it with the other rules we realize that φ satisfies much more interesting properties than the other rules. This support the idea of defining rules through canonical matrices.

Finally, we obtain two axiomatic characterizations of φ . The second characterization says that φ is the only rule satisfying strong cost monotonicity, population monotonicity, and equal sharing of extra costs. We believe that this is a nice characterization because we use properties based in clear economic principles.

In the next theorem we prove that the new rule φ satisfies all the properties mentioned in the paper.

Theorem 1. φ satisfies *CS*, *CM*, *SCM*, *PM*, *CON*, *POS*, *SEP*, *ESEC*, *IIT*, and *EC*.

Proof. See the Appendix.

Remark 4. Sprumont (1990) introduced the Population Monotonic Allocation Schemes (*PMAS*) for *TU* games. Let (N_0, C) be a *cstp* and let C^* be its canonical matrix. By Proposition 3 (c), v_{C^*} is concave and hence it is possible to find a *PMAS* for v_{C^*} . Since $C \geq C^*$, $v_C(S) \geq v_{C^*}(S)$ for all $S \subset N$ and hence, the *PMAS* of v_{C^*} are also *PMAS* in v_C . Since φ satisfies *PM* we conclude that $\{\varphi(S_0, C)\}_{S \subset N}$ is a *PMAS* of v .

In the next table we summarize the properties satisfied by the rules mentioned in the paper (in the Appendix we prove that this table is correct).

	B	K	DK	φ
CS	YES	no	YES	YES
CM	no	YES	YES	YES
SCM	no	no	no	YES
PM	no	no	no	YES
CON	no	YES	no	YES
POS	YES	no	YES	YES
SEP	YES	no	YES	YES
$ESEC$	YES	YES	no	YES
IIT	no	no	no	YES
EC	no	no	no	YES

This table shows clearly that φ satisfies much more interesting properties than the other rules mentioned in the paper.

This table together with Proposition 5 (b) suggest that we must take a lot of care when we define rules which do not depend only on the canonical form. K and B use information which is not contained in the canonical form. Nevertheless this table suggests that they use this information in a wrong way because they lose many properties. A similar argument can be applied to DK .

Our feeling is that the canonical form is a good environment for defining rules in $cstp$.

Remark 5. A natural subclass of $cstp$ is the one which arises when we assume that for any $i, j \in N_0$, c_{ij} , the direct cost of connecting i and j , is not larger than the cost of connecting i and j indirectly through other agents. Formally, for all $i, j \in N_0$,

$$c_{ij} = \min_{g \in G_{ij}} c(N_0, C, g)$$

where G_{ij} is the set of paths connecting i and j .

It is not difficult to check that in this subclass of problems the only change in the table is that now K satisfies POS .

In the next theorem we give two characterizations of φ .

Theorem 2.(a) φ is the only rule satisfying IIT , SEP , and $ESEC$.

(b) φ is the only rule satisfying *EC*.

Proof. See the Appendix.

Since *PM* implies *SEP*, *SCM* implies *IIT*, and φ satisfies *PM* and *SCM* next corollary is a trivial consequence of Theorem 2.

Corollary 1. φ is the only rule satisfying *SCM*, *PM*, and *ESEC*.

Even though Corollary 1 is a trivial consequence of Theorem 3 (a) we state it explicitly because we believe that *PM* and *SCM* are more appealing properties than *SEP* and *IIT*.

Remark 6. Corollary 1 and Theorem 2 (a) are tight characterization results (the proof is in the Appendix).

6 Other approaches to the problem

In this section we present four alternative definitions for φ . The first one is, of course, the definition of the *P-value* given by Feltkamp *et al* (1994a). In the second one we write φ as the Shapley value of a game. In the third we define it looking only to the problem (N_0, C) in an intuitive way. The fourth is made considering the costs as indivisible goods.

We believe that these results are very interesting because we arrive to it from very different approaches. In some way we can consider φ as a "focal point" of *cstp*.

First approach: The *P-value*.

Feltkamp *et al* (1994a) introduce a rule for *cstp* called Equal Remaining Obligations. Later, Branzei *et al* (2003) study it more and call it *P-value*, which is the name we use. As we say before when we start to study this problem we didn't know the paper of Feltkamp *et al* (1994a), which is an unpublished paper. In January 2004 we come across the paper of Branzei *et al* (2003), which was available in the web in December 2003. Through this paper we know the first one.

In this section we prove that the *P-value*, which has been defined in a completely different way than φ , coincides with φ .

Second approach: The Shapley value.

We already know that φ is the Shapley value of (N, v_{C^*}) . This approach is interesting because we prove that φ is the Shapley value of a game which is independent of the canonical form.

Given a *cstp* (N_0, C) we denote by (S_0, C^{+T}) the *cstp* obtained from (N_0, C) assuming that agents of S have to be connected and agents of T are already connected. This means that $c_{ij}^{+T} = c_{ij}$ for all $i, j \in S$ and $c_{i0}^{+T} = \min_{j \in T_0} c_{ij}$ for all $i \in S$.

Notice that if $S = N \setminus \{i\}$ and $T = \{i\}$ this definition coincides with $((N \setminus \{i\})_0, C^{+i})$, which has been defined before property of *EC*.

Now, we associate to each *cstp* (N_0, C) a *TU* game (N, v_C^+) where for each $S \subset N$,

$$v_C^+(S) = m(S_0, C^{+(N \setminus S)}).$$

Usually, we write v^+ instead of v_C^+ .

Given $S \subset N$, $v^+(S)$ is the minimal cost of connecting all agents of S to the source assuming that agents of $N \setminus S$ are already connected. Note that $v(S)$ is the minimal cost of connecting all agents of S to the source assuming that agents in $N \setminus S$ are not participating.

If we compute v and v^+ in Example 1 with $x = 100$ we obtain

S	v	v^+
$\{1\}$	10	2
$\{2\}$	100	2
$\{1,2\}$	12	12

This example shows that v and v^+ are very different. Nevertheless, for all *cstp* (N_0, C) we have that $v(N) = v^+(N) = m(N_0, C)$.

Definition 2. For all *cstp* (N_0, C) , we define

$$\varphi^1(N_0, C) = Sh(N, v^+).$$

In the next proposition we give some results about v^+ and φ^1 which will be useful to prove that φ^1 coincides with φ . Moreover, we also prove that φ^1 satisfies *EC*.

Proposition 6. (a) If C is a canonical matrix then, $v(S) + v^+(N \setminus S) = m(N_0, C)$ for all $S \subset N$.

(b) φ^1 satisfies *IIT*.

(c) φ^1 satisfies *EC*.

Proof. See the Appendix.

Third approach: Looking to the problem.

We now introduce a rule to *cstp* looking only to the problem. Assume that there is a unique *mcst* $t = \{(i^0, i)\}_{i \in N}$ in (N_0, C) . The problem that agents face is how to divide the cost induced by t among agents. We believe that a reasonable approach is to decide, for each arc of t , the part of the cost every agent has to pay. Notice that *B* and *DK* decide to assign (with different criteria) the cost of any arc to a single agent. We allow to divide the cost of an arc among several agents.

If we back to Example 1 with $x = 100$ we realize that the unique *mcst* is $t = \{(0, 1), (1, 2)\}$. We argue that both agents have rights over the cost of arc $(1, 2)$. In the case of agent 2 this is clear because he connects to the source through this arc. We also believe that agent 1 has rights over this cost. Agent 2 can connect through agent 1 only because agent 1 connects to the source. We believe that a fair rule must assign some of the benefits that agent 2 obtains by connecting through agent 1, which can be identified with the small cost of arc $(1, 2)$, to agent 1.

Since both agents are using the arc $(0, 1)$ for connecting to the source, we believe that any fair rule must assign this cost between both agents.

We then define a multi-step procedure to compute the part of the cost of any arc of t that every agent has to pay. The idea is the following. We ask agents by the cost of the arc of t they want to pay for. In order to take into account the structure of t , each agent must select an arc which he belongs to. We assume each agent chooses the arc with the lowest cost. If several agents select the same arc we divide this cost equally among the agents who selected it. Each agent pays the same proportion of the cost of the arc he selected. It is trivial to see that in the first step this proportion should be 0.5 or 1.

In the next steps we apply the same idea but now agents can also select other arcs. Agent i can select arc (j, k) if there is a path in t connecting i with j or k through arcs whose cost have been already paid.

In Example 1 each agent selects the arc $(1, 2)$. Since both agents select the same arc this is divided equally between both. Then, agents 1 and 2 pay half of the cost of arc $(1, 2)$. In step 2 both agents select the arc $(0, 1)$ and the cost is divided equally among them. Agent 2 selects $(0, 1)$ because 2 is

connected to 1 through arc $(1, 2)$, which has been paid in step 1.

If there are several *mcst* we select any of them. Later we will prove that the final allocation is independent of the selected *mcst*.

We now present this procedure in a formal way. Let t be a *mcst* in (N_0, C) .

Step 0. We consider:

$a^0(i) = \emptyset$ for all $i \in N$. In general $a^0(i)$ belongs to t .

$p^0 = 0$.

$\varrho^0(i, j) = 0$ for all $(i, j) \in t$.

$A^0 = \{(i, j) \in t \text{ such that } \varrho^0(i, j) < 1\}$. Notice that $A^0 = t$.

$P^0 = \{P_1^0, \dots, P_n^0\}$ is a partition of N such that $P_i^0 = \{i\}$ for all $i \in N$.

$f_i^0 = 0$ for all $i \in N$.

Assume we have defined Step r for all $r \leq s - 1$. We now define Step s .

In order to clarify this definition we will compute Step 1 of the process when applied to the following example.

Example 3. Let (N_0, C) be the *cstp* with *mcst* $t = \{(0, 1), (1, 2), (2, 3)\}$ and $c_{01} = 12$, $c_{12} = 4$, and $c_{23} = 6$. We do not specify the cost of the other arcs because it is not necessary for the procedure.

Given $P_q^{s-1} \in P^{s-1}$, agents of P_q^{s-1} select, among the non-paid arcs, an arc connecting them with agents outside P_q^{s-1} whose cost they want to pay. Formally, let $(i_q, j_q) \in A^{s-1}$ be such that $i_q \in P_q^{s-1}$, $j_q \in N_0 \setminus P_q^{s-1}$ and

$$c_{i_q j_q} = \min \{c_{ij} : (i, j) \in A^{s-1}, i \in P_q^{s-1}, \text{ and } j \in N_0 \setminus P_q^{s-1}\}.$$

If there exist many $(i_q, j_q) \in A^{s-1}$ we select any of them. We define $a^s(i) = (i_q, j_q)$ for all $i \in P_q^{s-1}$.

Example 3. For $P_1^0 = \{1\}$ there are two possibilities: $(0, 1)$ and $(1, 2)$. Then, $(i_1, j_1) = (1, 2)$. For $P_2^0 = \{2\}$ there are two possibilities: $(1, 2)$ and $(2, 3)$. Then, $(i_2, j_2) = (1, 2)$. For $P_3^0 = \{3\}$ there is only one possibility $(2, 3)$. Then, $(i_3, j_3) = (2, 3)$. Moreover, $a^1(1) = a^1(2) = (1, 2)$, and $a^1(3) = (2, 3)$.

We now compute p^s , the proportion (of the arc he selected) that any agent pays at this step. For each arc $(i, j) \in A^{s-1}$ we define

$$N_{ij}^s = \{k \in N : a^s(k) = (i, j)\} \text{ and}$$

$$p^s = \min_{(i,j) \in A^{s-1}} \left\{ \frac{1 - \varrho^{s-1}(i, j)}{|N_{ij}^s|} \right\}.$$

We define $\varrho^s(i, j) = \varrho^{s-1}(i, j) + |N_{ij}^s| p^s$ for each $(i, j) \in A^{s-1}$. Then, $\varrho^s(i, j) \leq 1$ for each $(i, j) \in A^{s-1}$ and there exists at least one $(i', j') \in A^{s-1}$ such that $\varrho^s(i', j') = 1$. Notice that $\varrho^s(i, j)$ denotes the proportion of the cost of arc (i, j) paid at the end of Step s .

Example 3. $N_{01}^1 = \emptyset$, $N_{12}^1 = \{1, 2\}$, and $N_{23}^1 = \{3\}$. Then, $p^1 = \frac{1}{2}$ (we take $\frac{1}{0} = +\infty$). $\varrho^1(0, 1) = 0$, $\varrho^1(1, 2) = 1$, and $\varrho^1(2, 3) = \frac{1}{2}$.

We now compute the arcs whose cost has not been paid.

$$A^s = \{(i, j) \in A^{s-1} \text{ such that } \varrho^s(i, j) < 1\}.$$

We define P^s joining elements of P^{s-1} in the following way. Two elements P_q^{s-1} and $P_{q'}^{s-1}$ of P^{s-1} are in the same element of P^s if $(i_{q'}, j_{q'}) = (i_q, j_q)$ and $\varrho^s(i_q, j_q) = 1$.

Example 3. $A^1 = \{(0, 1), (2, 3)\}$ and $P^1 = \{\{1, 2\}, \{3\}\}$

Finally, for each $i \in P_q^{s-1}$ we compute the cost that agent i pays in Step s .

$$f_i^s = p^s c_{i_q j_q}.$$

Example 3. $f_1^1 = 2$, $f_2^1 = 2$, and $f_3^1 = 3$.

This process ends when $A^s = \emptyset$. Since $A^0 = t$, $A^s \subset A^{s-1}$ and $A^s \neq A^{s-1}$, this process ends in a finite number of steps (at most n), say γ .

Since each tree has n arcs it is trivial to see that $\sum_{s=1}^{\gamma} p^s = 1$.

Definition 3. For all *cstp* (N_0, C) and $i \in N$ we define $\varphi_i^3(N_0, C) = \sum_{s=1}^{\gamma} f_i^s$.

We must prove that φ^3 is well defined (it does not depend on things such as the tree t , the arc (i_q, j_q) ...). We will do it later.

Remark 7¹. Feltkamp, Tijs, and Muto (1994b) introduce the decentralized rule for generic *cstp*. A *cstp* is generic if different arcs have different costs (without violating that $c_{ij} = c_{ji}$ for all $i, j \in N_0$). The decentralized rule is defined with a procedure similar to our procedure defining φ^3 . At each step each agent select a non paid arc with the same criteria that with our procedure.

The main difference between both procedures is the following. In the decentralized rule, all the cost of each arc selected is divided between the agents who select it. In our procedure it is possible that only a part of this cost is divided. We now compute the decentralized rule in Example 3.

Step 1. Agents 1 and 2 select arc $(1, 2)$. Agent 3 selects arc $(2, 3)$. Then, the cost of arc $(1, 2)$ is divided equally between agents 1 and 2. The cost of arc $(2, 3)$ is paid by agent 3. Agent 3 has not more obligations.

Step 2. Agents 1 and 2 select arc $(0, 1)$ and divided this cost equally among them.

Thus, agents 1 and 2 pay $\frac{1}{2}4 + \frac{1}{2}12 = 8$, whereas agent 3 pays 6.

Fourth approach: Indivisible costs.

To clarify the motivation of this approach we assume that there is a unique *mcst* in (N_0, C) . We can think in B and DK as rules assigning indivisible goods (cost of the arcs) because the cost of any arc of t is paid only by one agent. Of course, the indivisible goods are not really indivisible.

Given $\pi \in \Pi_N$ assume that the agents connect sequentially in the order given by π to one of the cheapest available options. Following this procedure we can associate a tree (not necessarily unique) to any permutation. We say that the tree t^π is induced by the permutation $\pi \in \Pi_N$ if:

$$t^\pi = \left\{ \begin{array}{l} (i^\pi, i) \text{ such that } i \in N, i^\pi \in Pre(i, \pi) \cup \{0\} \\ \text{and } c_{i^\pi i} \leq c_{ij} \text{ for all } j \in Pre(i, \pi) \cup \{0\} \end{array} \right\}.$$

Let us suppose that any agent pays his connection cost in accordance to t^π , *i.e.* agent i pays $c_{i^\pi i}$ for all $i \in N$. It is self-evident that if t^π and t'^π are two different trees induced by $\pi \in \Pi_N$, then agents pay the same under t^π as under t'^π . Thus it makes sense to define the cost allocation induced by the permutation $\pi \in \Pi_N$ as the vector $c^\pi = \{c_{i^\pi i}\}_{i \in N}$.

¹This remark is based on a suggestion of Stef Tijs.

Assume that $\pi \in \Pi_N$ satisfies $c(N_0, C, t^\pi) = m(N_0, C)$. Then, t^π is a *mcst* and $B(N_0, C)$ and $DK(N_0, C)$ are obtained from c^π . It is trivial to see that $B_i(N_0, C) = c_{i\pi_i}$ and $DK_i(N_0, C) = c_{i_j^\pi i_j}$ for all $i \in N$. Then, B and DK could be very unfair (see, for instance Example 1 with $x > 0$).

A classical way for recovering fairness from an allocation depending on the permutations is taking the average over the set of all permutations. In *cstp* problems this approach is incompatible with efficiency because for some $\pi \in \Pi_N$, $c(N_0, C, t^\pi) > m(N_0, C)$.

Nevertheless, if C is a canonical matrix, for all $\pi \in \Pi_N$, $c(N_0, C, t^\pi) = m(N_0, C)$ (we will prove this statement later). Then, in canonical matrices it is possible to generate an efficient and fair allocation taking the average over the permutations.

We then define the rule φ^3 as follows.

Definition 4. For all *cstp* (N_0, C) and $i \in N$ we define

$$\varphi_i^3(N_0, C) = \frac{1}{n!} \sum_{\pi \in \Pi_N} c_{i\pi_i}^*$$

where $c_{i\pi_i}^*$ is the connection cost among i and i^π in the canonical form C^* associated to C .

In this approach the canonical form appears in a natural way. Nevertheless, this approach is different from the original definition of φ .

We now prove that all the rules defined in this section coincide with φ .

Theorem 3. $P = \varphi^1 = \varphi^2 = \varphi^3 = \varphi$.

Proof. See the Appendix.

In the next corollary, which is a trivial consequence of Theorem 3 and Proposition 6 (c), we prove that φ satisfies *EC*.

Corollary 2. φ satisfies *EC*.

We believe that Theorem 3 gives more support to this rule because we can justify it from different approaches to the problem. This rule is in some sense a focal point of the problem.

7 Appendix

In this section we prove the results stated in the paper.

7.1 Proof of Proposition 1

We first prove that if C satisfies A1 and A2 then C is canonical. By simplicity we assume that $i_p = p$ for all $p = 0, 1, \dots, n$. This means that $t = \{(i-1, i)\}_{i=1}^n$.

Take $i, j \in N_0$ and C' such that $c'_{ij} < c_{ij}$ and $c'_{kl} = c_{kl}$ otherwise. We need to prove that $m(N_0, C') < m(N_0, C)$. It is enough to prove that there exists a tree t such that $c(N_0, C', t) < m(N_0, C)$.

Assume without loss of generality that $i < j$. By (A2), $c_{ij} = c_{(k-1)k}$ where $i < k \leq j$. We define $t' = (t \setminus \{(k-1, k)\}) \cup \{(i, j)\}$. Then, t' is a tree and

$$\begin{aligned} c(N_0, C', t') &= m(N_0, C) - c_{(k-1)k} + c'_{ij} \\ &= m(N_0, C) - c_{ij} + c'_{ij} \\ &< m(N_0, C). \end{aligned}$$

We now prove that if C is canonical then C satisfies A1 and A2.

Let (N_0, C) be a *cstp* and $t = \{(i^0, i)\}_{i \in N}$ a *mcst* in (N_0, C) . If $(j, k) \notin t$ there exists a unique path g in t connecting j and k . Take $(l, l^0) \in g$ such that $c_{l^0 l} = \max_{(i^0, i) \in g} \{c_{i^0 i}\}$. We now prove two claims.

Claim 1. $c_{jk} \geq c_{l^0 l}$.

Assume that $c_{jk} < c_{l^0 l}$. Take $t' = (t \setminus \{(l^0, l)\}) \cup \{(j, k)\}$. Then, t' is a tree and

$$c(N_0, C, t') = c(N_0, C, t) - c_{l^0 l} + c_{jk} < c(N_0, C, t)$$

which is a contradiction because t is a *mcst* in (N_0, C) .

Notice that Claim 1 holds also when C is not canonical.

Claim 2. $c_{jk} \leq c_{l^0 l}$.

Assume that $c_{jk} > c_{l^0 l}$. By Remark 2, we can find a *mcst* t' such that $(j, k) \in t'$. Let T_j be the set of nodes in N_0 which are connected to j in $t' \setminus \{(j, k)\}$. We can define T_k in a similar way. Notice that $j \in T_j$ and

$k \in T_k$. Since t' is a tree we can find $(p^0, p) \in g$ such that $(p^0, p) \notin t'$ and $t'' = (t' \setminus \{(j, k)\}) \cup \{(p^0, p)\}$ is a tree. Then,

$$c(N_0, C, t'') = c(N_0, C, t') - c_{jk} + c_{p^0 p} < c(N_0, C, t')$$

which is a contradiction because t' is a *mcst* in (N_0, C) .

We proceed by induction on n (the number of agents). If $n = 1$ the result is trivial. Assume that the result holds for $n \leq s$ and we prove it when $n = s + 1$.

We first prove that we can find a *mcst* satisfying (A1).

Let t' be a *mcst* obtained following Prim's algorithm. Then, t' can be obtained assuming that agents connect sequentially in some order $\pi \in \Pi_N$. By simplicity we take $\pi = \{1, 2, \dots, n\}$.

Because of the definition of Prim's algorithm we know that $t'_{N \setminus \{n\}}$ is a *mcst* in $((N \setminus \{n\})_0, C)$. Since $N \setminus \{n\}$ has $n - 1$ agents, by induction hypothesis, we can find $t^1 = \{(i_{p-1}, i_p)\}_{p=1}^{n-1}$ satisfying (A1) and (A2). Moreover,

$$c((N \setminus \{n\})_0, C, t'_{N \setminus \{n\}}) = c((N \setminus \{n\})_0, C, t^1).$$

Let $i_q \in \{0, 1, \dots, n - 1\}$ be the agent to whom n connects according with t' . We take $t^2 = t^1 \cup (i_q, n)$. It is easy to see that $c(N_0, C, t^2) = c(N_0, C, t)$. Thus, t^2 is a *mcst* in (N_0, C) .

For all $p = q + 1, \dots, n - 1$, $g_p = \{(i_q, n), \{(i_{r-1}, i_r)\}_{r=q+1}^p\}$ is the only path in t^2 connecting n and i_p . By claims 1 and 2, $c_{i_p n} = \max \left\{ \max_{r=q+1, \dots, p} \{c_{i_{r-1} i_r}\}, c_{i_q n} \right\}$.

Assume that there exist $r \in \{q + 1, \dots, n - 1\}$ such that $c_{i_{p-1} i_p} \leq c_{i_q n}$ for all $p = q + 1, \dots, r$ and $c_{i_q n} \leq c_{i_r i_{r+1}}$. It is trivial to see that the tree

$$t = \{(i_{p-1}, i_p)\}_{p=1}^r \cup (i_r, n) \cup (i_{r+1}, n) \cup \{(i_{p-1}, i_p)\}_{p=r+2}^{n-1}$$

is a *mcst* satisfying (A1) because $c_{i_r n} = c_{i_q n}$ and $c_{i_{r+1} n} = c_{i_r i_{r+1}}$.

Assume that $c_{i_{p-1} i_p} \leq c_{i_q n}$ for all $p = q + 1, \dots, n - 1$. Then, the tree

$$t = \{(i_{p-1}, i_p)\}_{p=1}^{n-1} \cup (i_q, n)$$

is a *mcst* satisfying (A1) because $c_{i_{n-1} n} = c_{i_q n}$.

We have proved that we can find a *mcst* $t = \{(i_{p-1}, i_p)\}_{p=1}^n$ satisfying (A1). We now prove that t also satisfies (A2).

For all $i_p, i_q \in N$, $p < q$ we have that $g = \{(i_{r-1}, i_r)\}_{r=p+1}^q$ is the unique path in t connecting i_p and i_q . By claims 1 and 2 we conclude that t also satisfies (A2).

7.2 Proof of Proposition 2

We prove several claims.

Claim 1. For all $p = 1, \dots, n$,

$$c_{i_p^0 i_p} = \min_{l \in S^{p-1}, k \in N_0 \setminus S^{p-1}} \{c_{lk}^{p-1}\}.$$

Because of the definition of C^{p-1} , $c_{lk}^{p-1} = c_{lk}$ when $l \in S^{p-1}$ and $k \in N_0 \setminus S^{p-1}$. Since $i_p^0 \in S^{p-1}$ and $i_p \in N_0 \setminus S^{p-1}$ we conclude that $c_{i_p^0 i_p} \geq \min_{l \in S^{p-1}, k \in N_0 \setminus S^{p-1}} \{c_{kl}\}$.

Assume that $c_{i_p^0 i_p} > \min_{l \in S^{p-1}, k \in N_0 \setminus S^{p-1}} \{c_{lk}\} = c_{l_p k_p}$. We take $t' = (t^0 \setminus \{(i_p^0, i_p)\}) \cup \{(l_p, k_p)\}$. Then, t' is a tree and

$$c(N_0, C, t') = c(N_0, C, t^0) - c_{i_p^0 i_p} + c_{l_p k_p} < c(N_0, C, t^0)$$

which is a contradiction because t^0 is a *mcst* in (N_0, C) .

Claim 2. For all $p = 1, \dots, n$, $C^{p-1} \geq C^p$ and C^{p-1} and C^p are directly equivalent.

We prove that $C^{p-1} \geq C^p$. By definition $c_{jk}^p = c_{jk}^{p-1}$ if $\{j, k\} \neq \{i_{p-1}, i_p\}$ it is enough to prove that $c_{i_{p-1} i_p}^{p-1} \geq c_{i_{p-1} i_p}^p$. Since $i_{p-1} \in S^{p-1}$, by Claim 1, $c_{i_{p-1} i_p}^{p-1} \geq c_{i_p^0 i_p} = c_{i_{p-1} i_p}^p$.

Assume that t' is a tree obtained through Prim's algorithm in (N_0, C^{p-1}) . By Claim 1 it is straightforward to prove that t' can be obtained also through Prim's algorithm in (N_0, C^p) . Now is easy to conclude that C^{p-1} and C^p are directly equivalent.

Claim 3. $C^n \geq C^*$.

We must prove that $c_{i_p i_q}^n \geq c_{i_p i_q}^* = \max_{p < r \leq q} \{c_{i_{r-1} i_r}^n\}$ for all $i_p, i_q \in N_0$, $p < q$. Suppose not. Then, there exist $i_p, i_q, i_{r-1}, i_r \in N_0$, $p < r \leq q$ such that $c_{i_{r-1} i_r}^n > c_{i_p i_q}^n$.

As a consequence of Prim's algorithm, $t = \{(i_{p-1}, i_p)\}_{p=1}^n$ is a *mcst* of (N_0, C^n) . We consider the tree $t' = (t \setminus \{(i_{r-1}, i_r)\}) \cup \{(i_p, i_q)\}$. Then,

$$c(N_0, C^n, t') = c(N_0, C^n, t) - c_{i_{r-1}i_r}^n + c_{i_p i_q}^n < c(N_0, C^n, t)$$

which is a contradiction because t is a *mcst* of (N_0, C^n) .

Claim 4. C^n and C^* are directly equivalent.

In the proof of Claim 3 we argue that $t = \{(i_{p-1}, i_p)\}_{p=1}^n$ is a *mcst* in (N_0, C^n) .

We now apply Prim's algorithm to (N_0, C^*) . Stage 1. We must select an arc $(0, i)$ such that $c_{0i}^* = \min_{j \in N} \{c_{0j}^*\}$. We know that for all $p = 1, \dots, n$, $q = 1, \dots, n$ and $p > q$, $c_{0i_p}^* \geq c_{0i_q}^*$. Then, in Stage 1 we can select the arc $(0, i_1)$.

Repeating this argument we can prove that in each Stage p , $p = 1, \dots, n$ we can select the arc (i_{p-1}, i_p) .

Thus, t is a *mcst* in (N_0, C^*) and hence, C^n and C^* are directly equivalent.

Claim 5. C^* is a canonical matrix, $C \geq C^*$, C and C^* are equivalent, and C^* is a canonical matrix associated to C .

By Proposition 1 C^* is a canonical matrix.

By Claim 2, $C = C^0 \geq C^1 \geq \dots \geq C^n$. By Claim 3, $C^n \geq C^*$. Then, $C \geq C^*$.

Since C^{p-1} and C^p are directly equivalent for all $p = 1, \dots, n$ (Claim 2) and $C^0 = C$, we conclude that C and C^n are equivalent. By Claim 4, C^n and C^* are equivalent. Then, C and C^* are equivalent.

Since C^* is a canonical matrix, $C \geq C^*$, and $m(N_0, C) = m(N_0, C^*)$ (C and C^* are equivalent) we conclude that C^* is a canonical matrix associated to C .

Claim 6. C^* is the unique canonical matrix associated to C .

Since C^* is a canonical matrix associated to C we must prove that if C' is another canonical matrix associated to C then, $C' = C^*$. Because of the definition of canonical matrices it is enough to prove that $C' \geq C^*$.

Since C^* and C' are canonical matrices associated to C and t^0 is a *mcst* in (N_0, C) , t^0 is also a *mcst* in (N_0, C^*) and (N_0, C') . Moreover, $c'_{i^0 i} = c_{i^0 i}^* = c_{i^0 i}$ for all $i \in N$.

Let t be the tree obtained in Stage $n + 1$ of the algorithm. Assume, without loss of generality, that $t = \{(i - 1, i)\}_{i=1}^n$.

We prove that given $i, j \in N_0$, $c'_{ji} \geq c^*_{ji}$. Assume, without loss of generality, that $i > j$. Since $c_{k^0k} = c^*_{(k-1)k}$ (because of the definition of t),

$$c^*_{ji} = \max_{j < k \leq i} \{c^*_{(k-1)k}\} = \max_{j < k \leq i} \{c_{k^0k}\}.$$

Let $k \in N$ be such that $j < k \leq i$. We consider two cases:

1. k is in the unique path in t^0 from i to 0. Then, $t' = t^0 \setminus \{(k^0, k)\} \cup \{(j, i)\}$ is a tree. Since t^0 is a *mcst* in (N_0, C') , Thus,

$$c(N_0, C', t^0) \leq c(N_0, C', t') = c(N_0, C', t^0) - c_{k^0k} + c'_{ji}.$$

which means that $c'_{ji} \geq c_{k^0k}$.

2. k is not in the unique path in t^0 from j to 0. Let l be such that $k < l \leq j$, l is in the unique path in t^0 from j to 0, and for all $r \in \{k, k+1, \dots, l-1\}$, r is not in the unique path in t^0 from j to 0. Because of the definition of the algorithm we know that $\{l^0, k^0\} \subset S^{k-1}$, $\{l, k\} \subset N_0 \setminus S^{k-1}$, and

$$c'_{k^0k} = c_{k^0k} = \min_{p \in S^{k-1}, q \in N_0 \setminus S^{k-1}} \{c_{pq}\} \leq c_{l^0l} = c'_{l^0l}.$$

By case 1, $c'_{ji} \geq c_{l^0l}$ and then, $c'_{ji} \geq c_{k^0k}$.

Now $c'_{ji} \geq \max_{i < k \leq j} \{c_{k^0k}\} = c^*_{ji}$ is a trivial consequence of cases 1 and 2.

7.3 Proof of Proposition 3

(a) We compute t' following Prim's algorithm. Since t satisfies (A2) and $i_{p-1} \leq i_p$ for all $p = 1, \dots, s$ we conclude that $c_{0i_1} \leq c_{0i_p}$ for all $p = 1, \dots, |S|$. Then, the arc $(0, i_1)$ could be the first arc in Prim's algorithm.

Since t satisfies (A2) we conclude that $c_{0i_2} \leq c_{0i_p}$ and $c_{i_1i_2} \leq c_{i_1i_p}$ for all $p = 2, \dots, |S|$. Then, $\min\{c_{0i_2}, c_{i_1i_2}\} \leq \min\{c_{0i_p}, c_{i_1i_p}\}$ for all $p = 2, \dots, |S|$. By (A2), $c_{i_1i_2} \leq c_{0i_2}$. Thus, $c_{i_1i_2} = \min\{c_{0i_2}, c_{i_1i_2}\}$ and, hence, we can choose the arc (i_1, i_2) as the second arc in Prim's algorithm.

If we continue with this process we obtain that $t' = \{(i_{p-1}, i_p)\}_{p=1}^{|S|}$ is a tree obtained following Prim's algorithm. This means that t' is a *mcs*t in (S_0, C) and hence,

$$v(S) = m(S_0, C) = c(S_0, C, t') = \sum_{p=1}^{|S|} c_{i_{p-1}i_p}.$$

(b) Assume first that $i_q \neq i_{|S|}$. By (a),

$$\begin{aligned} v(S) - v(S \setminus \{i_q\}) &= \sum_{p=1}^{|S|} c_{i_{p-1}i_p} - \left(\sum_{p=1}^{q-1} c_{i_{p-1}i_p} + c_{i_{q-1}i_{q+1}} + \sum_{p=q+1}^{|S|} c_{i_{p-1}i_p} \right) \\ &= c_{i_{q-1}i_q} + c_{i_qi_{q+1}} - c_{i_{q-1}i_{q+1}}. \end{aligned}$$

Since t satisfies (A2) we conclude that $c_{i_{q-1}i_{q+1}} = \max\{c_{i_{q-1}i_q}, c_{i_qi_{q+1}}\}$. Then,

$$v(S) - v(S \setminus \{i_q\}) = \min\{c_{i_{q-1}i_q}, c_{i_qi_{q+1}}\}.$$

Assume now that $i_q = i_{|S|}$. By (a),

$$v(S) - v(S \setminus \{i_q\}) = \sum_{p=1}^{|S|} c_{i_{p-1}i_p} - \sum_{p=1}^{|S|-1} c_{i_{p-1}i_p} = c_{i_{|S|-1}i_{|S|}}.$$

(c) Let S' be such that $S \subset S' \subset N$. Assume that $S' = \{i'_1, \dots, i'_{s'}\}$ and $i'_{p-1} \leq i'_p$ for all $p = 1, \dots, |S'|$. Notice that $|S'| \geq |S|$ because $S \subset S'$. We prove that for all $i \in S$,

$$v(S) - v(S \setminus \{i\}) \geq v(S') - v(S' \setminus \{i\}).$$

We consider three cases:

- $i \neq i_{|S|}$. Then, $i = i_q = i'_{q'}$ and $i'_{q'} \neq i'_{|S'|}$. By (b),

$$\begin{aligned} v(S) - v(S \setminus \{i\}) &= \min\{c_{i_{q-1}i}, c_{ii_{q+1}}\} \\ v(S') - v(S' \setminus \{i\}) &= \min\{c_{i'_{q'-1}i}, c_{ii'_{q'+1}}\}. \end{aligned}$$

Since $S \subset S'$ we conclude that $i_{q-1} \leq i'_{q'-1}$ and $i'_{q'+1} \leq i_{q+1}$. Then, $c_{i_{q-1}i} \geq c_{i'_{q'-1}i}$ and $c_{ii_{q+1}} \geq c_{ii'_{q'+1}}$ because t satisfies (A2). Thus,

$$v(S) - v(S \setminus \{i\}) \geq v(S') - v(S' \setminus \{i\}).$$

- $i = i_{|S|}$ and $i \neq i'_{|S'|}$. Then $i = i_{|S|} = i'_{q'} < i'_{|S'|}$. By (b),

$$\begin{aligned} v(S) - v(S \setminus \{i\}) &= c_{i_{|S|-1}i} \\ v(S') - v(S' \setminus \{i\}) &= \min \left\{ c_{i'_{q'-1}i}, c_{ii'_{q'+1}} \right\}. \end{aligned}$$

Since $S \subset S'$ we conclude that $i_{|S|-1} \leq i'_{q'-1}$. Then, $c_{i_{|S|-1}i} \geq c_{i'_{q'-1}i}$ because t satisfies (A2). Thus,

$$v(S) - v(S \setminus \{i\}) \geq v(S') - v(S' \setminus \{i\}).$$

- $i = i_{|S|} = i'_{|S'|}$. By (b),

$$\begin{aligned} v(S) - v(S \setminus \{i\}) &= c_{i_{|S|-1}i} \\ v(S') - v(S' \setminus \{i\}) &= c_{i'_{|S'|-1}i}. \end{aligned}$$

Since $S \subset S'$ we conclude that $i_{|S|-1} \leq i'_{|S'|-1}$. Then, $c_{i_{|S|-1}i} \geq c_{i'_{|S'|-1}i}$ because t satisfies (A2).

7.4 Proof of Proposition 4

Let $t = \{(i_{p-1}, i_p)\}_{p=1}^n$ be the tree associated to C satisfying (A1).

We know that for all $i \in N$,

$$\begin{aligned} B_i(N_0, C) &= \frac{1}{n!} \sum_{\pi \in \Pi_N} B_i^\pi(N_0, C) \text{ and} \\ K_i(N_0, C) &= \frac{1}{n!} \sum_{\pi \in \Pi_N} (v_C(\text{Pre}(i, \pi) \cup \{i\}) - v_C(\text{Pre}(i, \pi))). \end{aligned}$$

We prove that B coincides with K using an induction argument over the set of agents. If $n = 1$ it is clear that B coincides with K . Assume that $B(N_0, C) = K(N_0, C)$ when $n \leq p - 1$.

We prove it when $n = p$. We consider two cases.

Case 1. There exists an arc (i_{q-1}, i_q) with $q > 1$ such that $c_{i_{q-1}i_q} \geq c_{0i_1}$.

We take $S = \{i_1, \dots, i_{q'-1}\}$ where q' satisfies that $c_{i_{q'-1}i_{q'}} \geq c_{0i_1}$ and $c_{i_{r-1}i_r} < c_{0i_1}$ for all $r < q'$.

By Proposition 3 (a) and condition (A2),

$$\begin{aligned}
m(N_0, C) &= \sum_{r=1}^n c_{i_{r-1}i_r} = \sum_{r=1}^{q'-1} c_{i_{r-1}i_r} + \sum_{r=q'}^n c_{i_{r-1}i_r} \\
&= m(S_0, C) + c_{i_{q'-1}i_{q'}} + \sum_{r=q'+1}^n c_{i_{r-1}i_r} \\
&= m(S_0, C) + c_{0i_{q'}} + \sum_{r=q'+1}^n c_{i_{r-1}i_r} \\
&= m(S_0, C) + m((N \setminus S)_0, C).
\end{aligned}$$

Later, we will prove that B and K satisfy *SEP* without using this proposition. Then,

$$\begin{aligned}
B_i(N_0, C) &= \begin{cases} B_i(S_0, C) & \text{if } i \in S \\ B_i((N \setminus S)_0, C) & \text{if } i \in N \setminus S \end{cases} \quad \text{and} \\
K_i(N_0, C) &= \begin{cases} K_i(S_0, C) & \text{if } i \in S \\ K_i((N \setminus S)_0, C) & \text{if } i \in N \setminus S \end{cases}
\end{aligned}$$

Since S and $N \setminus S$ has at most $q - 1$ agents, by induction hypothesis we conclude that B coincides with K .

Case 2. $c_{0i_1} > c_{i_{p-1}i_p}$ for all $p = 2, \dots, n$.

Take $i \in N$. Then, $i = i_p$ for some $p = 1, \dots, n$. Since t satisfies (A2)

$$c_{0i} = \max_{r=1, \dots, p} \{c_{i_{r-1}i_r}\} = c_{0i_1}.$$

Given $i \in N$ we define the *cstp* (N_0^{-i}, C^i) where $N^{-i} = N \setminus \{i\}$ and for all $j, k \in N_0^{-i}$, $j \leq k$

$$c_{jk}^i = \begin{cases} c_{jk} & \text{if } j \neq 0 \\ c_{ik} & \text{if } j = 0. \end{cases}$$

Since C^i is symmetric we can compute c_{jk}^i when $j > k$.

Notice that (N_0^{-i}, C^i) is obtained from (N_0, C) by considering agent i as the source. We then write N_i^{-i} instead of N_0^{-i} .

We prove several claims.

Claim 1. C^i is a canonical matrix and $m(N_i^{-i}, C^i) = m(N_0, C) - c_{0i_1}$.

We first prove that $m(N_i^{-i}, C^i) = m(N_0, C) - c_{0i_1}$. We know that $i = i_q$ for some $q = 1, \dots, n$. Then, $t^i = t \setminus \{(0, i_1)\}$ is a tree in (N_0^{-i}, C^i) and $c(t^i, N_i^{-i}, C^i) = m(N_0, C) - c_{0i_1}$. Thus, $m(N_i^{-i}, C^i) \leq m(N_0, C) - c_{0i_1}$.

Assume that $m(N_i^{-i}, C^i) < m(N_0, C) - c_{0i_1}$. There exists a tree t' in (N_0^{-i}, C^i) satisfying that $c(t', N_i^{-i}, C^i) < c(t^i, N_i^{-i}, C^i)$. We define $t'' = t' \cup (0, i)$. Since t'' is a tree in (N_0, C)

$$\begin{aligned} c(t'', N_0, C) &= c_{0i} + c(t', N_i^{-i}, C^i) < c_{0i_1} + c(t^i, N_i^{-i}, C^i) \\ &= m(N_0, C) \end{aligned}$$

which is a contradiction. Then, $m(N_i^{-i}, C^i) = m(N_0, C) - c_{0i_1}$.

Using arguments similar to those used in the proof of $m(N_i^{-i}, C^i) < m(N_0, C) - c_{0i_1}$ we can prove that if we reduce the cost of an arc in (N_i^{-i}, C^i) then the *mcst* in (N_i^{-i}, C^i) is also reduced. Thus, C^i is a canonical matrix. This ends the proof of Claim 1.

Given $i \in N$ we define $\Pi^i = \{\pi \in \Pi_N \text{ such that } \pi(1) = i\}$, *i.e.* the orders in which agent i is the first.

Claim 2. For all $i \in N$ and $j \in N^{-i}$,

$$B_j(N_i^{-i}, C^i) = \frac{1}{(n-1)!} \sum_{\pi \in \Pi^i} B_j^\pi(N_0, C).$$

Take $j \in N^{-i}$. Since t satisfies (A2) and $c_{0i_1} > c_{i_{p-1}i_p}$ for all $p = 2, \dots, n$, we conclude that $c_{ji} < c_{j0}$. Because of the definition of Prim's algorithm, for all $\pi \in \Pi^i$,

$$B_j^\pi(N_0, C) = B_j^{\pi_{N^{-i}}}(N_i^{-i}, C^i).$$

Since $\pi \in \Pi^i$ if and only if $\pi_{N^{-i}} \in \Pi_{N^{-i}}$,

$$B_j(N_i^{-i}, C^i) = \frac{1}{(n-1)!} \sum_{\pi \in \Pi_{N^{-i}}} B_j^\pi(N_i^{-i}, C^i) = \frac{1}{(n-1)!} \sum_{\pi \in \Pi^i} B_j^\pi(N_0, C).$$

Claim 3. For all $i \in N$ and $j \in N^{-i}$,

$$K_j(N_i^{-i}, C^i) = \frac{1}{(n-1)!} \sum_{\pi \in \Pi^i} v_C(\text{Pre}(j, \pi) \cup \{j\}) - v_C(\text{Pre}(j, \pi)).$$

Using arguments similar to those used when we prove that $m(N_i^{-i}, C^i) = m(N_0, C) - c_{0i_1}$ it is easy to prove that if $i \in S \subset N$,

$$m(S_0, C) = m((S \setminus \{i\})_i, C^i) + c_{0i}.$$

Then, for all $\pi \in \Pi^i$ and $j \in N^{-i}$,

$$\begin{aligned} v_C(\text{Pre}(j, \pi) \cup \{j\}) - v_C(\text{Pre}(j, \pi)) &= \\ &= m((\text{Pre}(j, \pi) \cup \{j\})_0, C) - m(\text{Pre}(j, \pi)_0, C) \\ &= m((\text{Pre}(j, \pi) \cup \{j\} \setminus \{i\})_i, C^i) - m((\text{Pre}(j, \pi) \setminus \{i\})_i, C^i) \\ &= v_{C^i}(\text{Pre}(j, \pi_{N^{-i}}) \cup \{j\}) - v_{C^i}(\text{Pre}(j, \pi_{N^{-i}})). \end{aligned}$$

Since $\pi \in \Pi^i$ if and only if $\pi_{N^{-i}} \in \Pi_{N^{-i}}$,

$$\begin{aligned} K_j(N_i^{-i}, C^i) &= \frac{1}{(n-1)!} \sum_{\pi \in \Pi_{N^{-i}}} v_{C^i}(\text{Pre}(j, \pi) \cup \{j\}) - v_{C^i}(\text{Pre}(j, \pi)) \\ &= \frac{1}{(n-1)!} \sum_{\pi \in \Pi^i} v_C(\text{Pre}(j, \pi) \cup \{j\}) - v_C(\text{Pre}(j, \pi)). \end{aligned}$$

Claim 4. For all $i \in N$ and $\pi \in \Pi^i$,

$$B_i^\pi(N_0, C) = v_C(\text{Pre}(i, \pi) \cup \{i\}) - v_C(\text{Pre}(i, \pi)) = c_{0i_1}.$$

Take $i \in N$ and $\pi \in \Pi^i$.

We know that $c_{0j} = c_{0i_1}$ for all $j \in N$. Then, in Stage 1 of Prim's algorithm we can select any agent $j \in N$. Since $\pi(1) = i$, $B_i^\pi(N_0, C) = c_{0i} = c_{0i_1}$.

Moreover,

$$\begin{aligned} v_C(\text{Pre}(i, \pi) \cup \{i\}) - v_C(\text{Pre}(i, \pi)) &= v_C(\{i\}) - v_C(\emptyset) = v_C(\{i\}) \\ &= m(\{i\}_0, C) = c_{0i} = c_{0i_1}. \end{aligned}$$

Claim 5. For all $j \in N$, $B_j(N_0, C) = K_j(N_0, C)$.

Take $j \in N$.

By claims 2 and 4,

$$\begin{aligned}
B_j(N_0, C) &= \frac{1}{n!} \sum_{\pi \in \Pi_N} B_j^\pi(N_0, C) = \frac{1}{n!} \sum_{i \in N} \sum_{\pi \in \Pi^i} B_j^\pi(N_0, C) \\
&= \frac{1}{n!} \sum_{\pi \in \Pi^j} B_j^\pi(N_0, C) + \frac{1}{n} \sum_{i \in N \setminus \{j\}} \frac{1}{(n-1)!} \sum_{\pi \in \Pi^i} B_j^\pi(N_0, C) \\
&= \frac{1}{n!} \sum_{\pi \in \Pi^j} c_{0i_1} + \frac{1}{n} \sum_{i \in N \setminus \{j\}} B_j(N_i^{-i}, C^i) \\
&= \frac{1}{n} c_{0i_1} + \frac{1}{n} \sum_{i \in N \setminus \{j\}} B_j(N_i^{-i}, C^i).
\end{aligned}$$

By claims 3 and 4,

$$\begin{aligned}
K_j(N_0, C) &= \frac{1}{n!} \sum_{\pi \in \Pi_N} (v_C(\text{Pre}(j, \pi) \cup \{i\}) - v_C(\text{Pre}(j, \pi))) \\
&= \frac{1}{n!} \sum_{i \in N} \sum_{\pi \in \Pi^i} (v_C(\text{Pre}(j, \pi) \cup \{i\}) - v_C(\text{Pre}(j, \pi))) \\
&= \frac{1}{n!} \sum_{\pi \in \Pi^j} (v_C(\text{Pre}(j, \pi) \cup \{i\}) - v_C(\text{Pre}(j, \pi))) \\
&\quad + \frac{1}{n} \sum_{i \in N \setminus \{j\}} \frac{1}{(n-1)!} \sum_{\pi \in \Pi^i} (v_C(\text{Pre}(j, \pi) \cup \{i\}) - v_C(\text{Pre}(j, \pi))) \\
&= \frac{1}{n!} \sum_{\pi \in \Pi^j} c_{0i_1} + \frac{1}{n} \sum_{i \in N \setminus \{j\}} K_j(N_i^{-i}, C^i) \\
&= \frac{1}{n} c_{0i_1} + \frac{1}{n} \sum_{i \in N \setminus \{j\}} K_j(N_i^{-i}, C^i).
\end{aligned}$$

By induction hypothesis we know that $B_j(N_i^{-i}, C^i) = K_j(N_i^{-i}, C^i)$ for all $i, j \in N$. Then, $B_j(N_0, C) = K_j(N_0, C)$.

7.5 Proof of Proposition 5

(a) Assume that ψ satisfies *IIT*. By Proposition 2, C is equivalent to C^* . Then, $\psi(N_0, C) = \psi(N_0, C^*)$.

Assume that ψ depends only on the canonical form. Because of the definition of equivalent problems, it is enough to prove that, if C and C' are two directly equivalent problems, then $\psi(N_0, C) = \psi(N_0, C')$.

Assume that C and C' are directly equivalent problems. There exists a *mcst* $t = \{(i^0, i)\}_{i \in N}$ in (N_0, C) and (N_0, C') such that $c_{i^0 i} = c'_{i^0 i}$ for all $i \in N$. It is straightforward to prove, because of the definition of the algorithm, that $C^* = C'^*$. Since ψ depends only on the canonical form, $\psi(N_0, C) = \psi(N_0, C')$.

(b) If ψ satisfies *SCM*, $\psi(N_0, C) \geq \psi(N_0, C^*)$ because $C \geq C^*$ (Proposition 2). Since $m(N_0, C) = m(N_0, C^*)$, $\psi(N_0, C) = \psi(N_0, C^*)$. By (a) we conclude that ψ satisfies *IIT*.

7.6 Proof of Theorem 1

Let (N_0, C) be a *cstp* and (N_0, C^*) its canonical form. We know that $\varphi(N_0, C) = K(N_0, C^*) = Sh(N, v_{C^*})$. We now prove that φ satisfies the following properties:

Independence of irrelevant trees (IIT). It is a consequence of Proposition 5 (a).

Strong cost monotonicity (SCM). Let (N_0, C) and (N_0, C') be two *cstp* such that there exist $j, k \in N_0$ satisfying $c'_{jk} = c_{jk} + a$ where $a > 0$ and $c'_{il} = c_{il}$ otherwise. We must prove that $\varphi_i(N_0, C) \leq \varphi_i(N_0, C')$ for all $i \in N$.

Since $C \leq C'$ we conclude that $m(N_0, C) \leq m(N_0, C')$.

Assume first there exists a *mcst* t in (N_0, C) such that $(j, k) \notin t$. Then,

$$c(N_0, C', t) = c(N_0, C, t) = m(N_0, C).$$

Thus, $m(N_0, C) = m(N_0, C')$ and t is a *mcst* in (N_0, C') . Moreover, $c_{il} = c'_{il}$ for all $(i, l) \in t$. This means that (N_0, C) and (N_0, C') are directly equivalent. Then, $\varphi(N_0, C) = \varphi(N_0, C')$ because φ satisfies *IIT*.

Assume that for all *mcst* t in (N_0, C) , $(j, k) \in t$. Let G be the set of trees which do not contain the arc (j, k) , *i.e.*

$$G = \{g \in \mathcal{G}_0^N \mid g \text{ is a tree and } (j, k) \notin g\}.$$

Let $t^1 \in G$ be such that

$$c(N_0, C, t^1) = \min_{t \in G} \{c(N_0, C, t)\}.$$

Since $(j, k) \notin t^1$, we conclude that $c(N_0, C, t^1) = c(N_0, C', t^1)$.

We define $b = c(N_0, C, t^1) - m(N_0, C)$. Notice that $b \geq 0$ because t^1 is a tree in (N_0, C) .

We now distinguish two cases:

Case 1: $a \leq b$. Let $t^0 = \{(i^0, i)\}_{i \in N}$ be a *mcst* in (N_0, C) . We assume, without loss of generality, that $j^0 = k$. Then, $(k, j) \in t^0$ and t^0 is a *mcst* in (N_0, C') . We can assume, without loss of generality, that t , the tree obtained when we apply the algorithm to (N_0, C) , is $\{(i-1, i)\}_{i \in N}$.

Let $t' = \{(i_{p-1}, i_p)\}_{p=1}^n$ be the tree obtained when we apply the algorithm to (N_0, C') . It is trivial to see that we can choose t' satisfying that $i_p = p$ for all $p < j$. Moreover, t' is computed in such a way that $t' = t$ if it is possible.

We consider two subcases:

Subcase 1.1: $t = t'$.

We first prove that $C^* \leq C'^*$.

When $i \neq j$, $c_{(i-1)i}^* = c_{i^0 i} = c'_{i^0 i} = c'_{(i-1)i}$. Moreover, $c_{(j-1)j}^* = c_{j^0 j} < c_{j^0 j} + a = c'_{(j-1)j}$.

Let $i, l \in N_0$ such that $i < l$. Because of the definition of C^* and C'^* we have that $c_{il}^* = \max_{i < p \leq l} \{c_{(p-1)p}^*\}$ and $c_{il}'^* = \max_{i < p \leq l} \{c_{(p-1)p}'^*\}$. Then, $c_{il}^* \leq c_{il}'^*$.

We now prove that $\varphi(N_0, C) \leq \varphi(N_0, C')$.

We know that $\varphi(N_0, C) = Sh(N, v_{C^*})$ and $\varphi(N_0, C') = Sh(N, v_{C'^*})$.

Take $S = \{j_1, \dots, j_{|S|}\}$ such that $j_{p-1} < j_p$ for all $p = 2, \dots, |S|$ and $j_q \in S$. By Proposition 3 (b) we conclude that

$$v_{C^*}(S) - v_{C^*}(S \setminus \{j_q\}) \leq v_{C'^*}(S) - v_{C'^*}(S \setminus \{j_q\})$$

because $C^* \leq C'^*$.

Take $i \in N$. Then,

$$\begin{aligned} Sh_i(N, v_{C^*}) &= \frac{1}{n!} \sum_{\pi \in \Pi_N} (v_{C^*}(Pre(i, \pi) \cup \{i\}) - v_{C^*}(Pre(i, \pi))) \\ &\leq \frac{1}{n!} \sum_{\pi \in \Pi_N} (v_{C'^*}(Pre(i, \pi) \cup \{i\}) - v_{C'^*}(Pre(i, \pi))) \\ &= Sh_i(N, v_{C'^*}). \end{aligned}$$

Subcase 1.2: $t \neq t'$.

It is trivial to see that $c_{j^0j} \leq c_{i_{j-1}i_j} < c_{j^0j} + a = c'_{j^0j}$. This means that $j = i_q$ with $q > j$. Because of the definition of the algorithm we can take t' such that $c'_{i_{p-1}i_p} < c'_{i_{q-1}i_q}$ for all $p = j, \dots, q-1$. Then, $c'_{i_{q-1}i_q} = c'_{j^0j} = c_{j^0j} + a$.

Take $C'^{(j-1)} = C$. For all $p = j, \dots, q$ we define C'^p such that $c'_{j^0j} = c'^{(p-1)}_{j^0j} + \max \left\{ c_{i_p^0i_p} - c'^{(p-1)}_{j^0j}, 0 \right\}$ and $c'_{il} = c'^{(p-1)}_{il}$ otherwise. Let t'^p be the tree obtained when we apply the algorithm to (N_0, C'^p) .

Notice that

$$C' = C'^q \geq C'^{(q-1)} \geq \dots \geq C'^j \geq C'^{(j-1)} = C.$$

and t^0 is a *mcst* in (N_0, C'^p) for all $p = j, \dots, q$.

We prove that for all $p = j, \dots, q$ we can take $t'^{(p-1)}$ and t'^p such that $t'^{(p-1)} = t'^p$. We prove it when $p = j$, the other cases are similar.

We take $t'^{(j-1)} = t$. If we apply the algorithm to C'^j , for all $i = 1, \dots, j-1$, we can select the arc (i^0, i) .

Since $c'_{j^0j} = c_{j^0j}$ and $c_{j^0j} \leq c_{i_j^0i_j}$,

$$c'_{j^0j} = c'_{j^0j} + \max \left\{ c_{i_j^0i_j} - c'^{(j-1)}_{j^0j}, 0 \right\} = c_{i_j^0i_j}.$$

Then, in Stage j of the algorithm we can select the arc (j^0, j) . Now it is trivial to see that in Stage i , $i > j$, we can select the arc (i^0, i) . Thus, $t'^j = t$.

Take $p \in \{j, \dots, q\}$. Since $t'^{(p-1)} = t'^p$ and $(N_0, C'^{(p-1)})$ and (N_0, C'^p) are in subcase 1.1 we conclude that $\varphi(N_0, C'^p) \geq \varphi(N_0, C'^{(p-1)})$.

Thus,

$$\varphi(N_0, C') = \varphi(N_0, C'^q) \geq \dots \geq \varphi(N_0, C'^{(j-1)}) = \varphi(N_0, C).$$

Case 2: $a > b$. Then, t^1 is a *mcst* in (N_0, C') but t is not.

Let C'' such that $c''_{jk} = c_{jk} + b$ and $c''_{il} = c_{il}$ otherwise. Notice that $C \leq C'' \leq C'$.

We are then under the hypothesis of Case 1, and thus $\varphi_i(N_0, C) \leq \varphi_i(N_0, C'')$ for all $i \in N$.

We know that t^1 is a *mcst* of (N_0, C') and (N_0, C'') and $(k, j) \notin t^1$. Then, $c'_{il} = c''_{il}$ for all $(i, l) \in t^1$.

This means that (N_0, C') and (N_0, C'') are directly equivalent. Since φ satisfies *IIT* (Proposition 5 (a)) we conclude that $\varphi(N_0, C'') = \varphi(N_0, C')$. Thus, $\varphi(N_0, C) \leq \varphi_i(N_0, C')$.

Cost monotonicity (CM). Since φ satisfies SCM we conclude that φ also satisfies CM .

Population monotonicity (PM). We must prove that for all $cstp$ (N_0, C) , all $S \subset N$, and all $i \in S$

$$\varphi_i(N_0, C) \leq \varphi_i(S_0, C).$$

It is enough to prove it when $S = N \setminus \{j\}$ and $j \in N$.

We prove several claims.

Claim 1. Let (N_0, C) be such that there exists $j \in N$ satisfying: $c_{j0} = a$, $c_{ji} = b$ for all $i \in N \setminus \{j\}$, and $b > a > \max_{i,k \in N_0 \setminus \{j\}} \{c_{ik}\}$. Then,

$$\varphi_i(N_0, C) = \begin{cases} a & \text{if } i = j \\ \varphi_i((N \setminus \{j\})_0, C) & \text{otherwise.} \end{cases}$$

By simplicity we assume that $j = n$. Let t^0 be a $mcst$ in $((N \setminus \{n\})_0, C)$. Then, $t^0 = t'^0 \cup \{(n, 0)\}$ is a $mcst$ in (N_0, C) .

Let t' be the tree obtained when we apply the algorithm to $((N \setminus \{n\})_0, C)$ starting in t'^0 . By simplicity we assume that $t' = \{(i-1, i)\}_{i=1}^{n-1}$. It is trivial to see that $t = t' \cup \{(n-1, n)\}$ is the tree obtained when we apply the algorithm to (N_0, C) starting in t^0 . For all $i = 1, \dots, n-1$, the cost of the arc $(i-1, i)$ is the same in the canonical form associated to $((N \setminus \{n\})_0, C)$ and (N_0, C) . Moreover, the cost of arc $(n-1, n)$ in (N_0, C^*) is a .

Let v_{-n} be the characteristic function associated to the $cstp$ $((N \setminus \{n\})_0, C^*)$.

By Proposition 3 (a), $v_{C^+}(S) = v_{-n}(S \setminus \{n\}) + a$ if $n \in S$ and $v_{C^*}(S) = v_{-n}(S)$ if $n \notin S$.

Consider the TU games (N, w_1) and (N, w_2) where for all $S \subset N$

$$\begin{aligned} w_1(S) &= v_{-n}(S \setminus \{n\}) \text{ and} \\ w_2(S) &= \begin{cases} a & \text{if } n \in S \\ 0 & \text{if } n \notin S. \end{cases} \end{aligned}$$

It is trivial to see that

$$\begin{aligned} Sh_i(N, w_1) &= Sh_i(N \setminus \{n\}, v_{-n}) = \varphi_i((N \setminus \{j\})_0, C) \text{ if } i \neq n, \\ Sh_n(N, w_1) &= 0, \\ Sh_i(N, w_2) &= 0 \text{ if } i \neq n, \text{ and} \\ Sh_n(N, w_2) &= a. \end{aligned}$$

Since $v(S) = w_1(S) + w_2(S)$ for all $S \subset N$ and the Shapley value is additive on the characteristic function, Claim 1 holds.

Claim 2. Given $j \in N$, $\varphi_i(N_0, C) \leq \varphi_i((N \setminus \{j\})_0, C)$ for all $i \in N \setminus \{j\}$.

By simplicity assume that $j = n$. Let $a \in \mathbb{R}$ be such that $a > \max_{i,j \in N_0} \{c_{ij}\}$.

Take $b = a + 1$. Let C'^0 be such that $c'_{0n} = a$ and $c'_{ij} = c_{ij}$ otherwise, Let C'^1 be such that $c'_{1n} = b$ and $c'_{ij} = c'_{ij}^0$ otherwise. In general, for all $k = 2, \dots, n-1$ we define $c'_{kn} = b$ and $c'_{ij} = c'_{ij}^{k-1}$ otherwise.

Take $i \in N \setminus \{n\}$. Since φ satisfies *SCM*,

$$\varphi_i(N_0, C) \leq \varphi_i(N_0, C'^0) \leq \varphi_i(N_0, C'^1) \leq \dots \leq \varphi_i(N_0, C'^{n-1}).$$

By Claim 1, $\varphi_i(N_0, C'^{n-1}) = \varphi_i((N \setminus \{n\})_0, C'^{n-1})$. Since $c'_{kl} = c_{kl}$ for all $k, l \in (N \setminus \{n\})_0$ we conclude that

$$\varphi_i((N \setminus \{n\})_0, C'^{n-1}) = \varphi_i((N \setminus \{n\})_0, C).$$

Then, $\varphi_i(N_0, C) \leq \varphi_i((N \setminus \{n\})_0, C)$.

Continuity (CON). Since C^* is a continuous function of C , v_{C^*} is a continuous function of C^* , and Sh is a continuous function of v_{C^*} , φ is a continuous function of C .

Separability (SEP). We know that φ satisfies *PM*. Since *PM* implies *SEP*, the result holds.

Core selection (CS). We know that φ satisfies *PM*. Since *PM* implies *CS*, the result holds.

Equal Sharing of Extra Costs (ESEC). Let (N_0, C) and (N_0, C') be as in the definition of *ESEC*. Assume that t^0 is a *mcst* in (N_0, C) . Then, t^0 is also a *mcst* in (N_0, C') .

If $t = \{(i_{p-1}, i_p)\}_{p=1}^n$ is the tree given by the algorithm when we apply to (N_0, C) , t is also the tree given by the algorithm when we apply to (N_0, C') . Moreover, $c_{0i_1}^* = c_0$, $c_{0i_1}' = c_0'$, and $c_{i_{p-1}i_p}^* = c_{i_{p-1}i_p}'$ for all $p = 2, \dots, n$.

By Proposition 3 (a), $v_{C'^*}(S) = v_{C^*}(S) + (c_0' - c_0)$ for all $S \subset N$.

Then, for all $i \in N$

$$\begin{aligned} \varphi_i(N_0, C') &= Sh_i(N, v_{C'^*}) = Sh_i(N, v_{C^*}) + \frac{c_0' - c_0}{n} \\ &= \varphi_i(N_0, C) + \frac{c_0' - c_0}{n}. \end{aligned}$$

Positivity (POS). Given $S \subset N$, by Proposition 3 (b), $v_{C^*}(S) - v_{C^*}(S \setminus \{i\}) \geq 0$ for all $i \in N$. Then, $\varphi_i(N_0, C) = Sh_i(N, v_{C^*}) \geq 0$.

Equal contributions (EC). We will prove it in Corollary 2.

7.7 Proof of the table

About Bird's rule B .

- *CS*. YES. See Bird (1976).
- *CM*. NO. See Dutta and Kar (2002).
- *SCM*. NO because B does not satisfy *CM*.
- *PM*. NO. Let (N_0, C) be such that $N = \{1, 2, 3\}$, $c_{01} = 30$, $c_{02} = 100$, $c_{03} = 10$, $c_{12} = 10$, $c_{13} = 40$, and $c_{23} = 20$.
Then, $B_2(N_0, C) = 20$ but $B_2(\{1, 2\}_0, C) = 10$.
- *CON*. NO. See Example 1.
- *POS*. YES. It is trivial.
- *SEP*. YES. Set $S \subset N$ be such that $m(N_0, C) = m(S_0, C) + m((N \setminus S)_0, C)$. Applying Prim's algorithm it is trivial to see that $B_i^\pi(N_0, C) = B_i^{\pi_S}(S_0, C)$ when $i \in S$ and $B_i^\pi(N_0, C) = B_i^{\pi_{N \setminus S}}((N \setminus S)_0, C)$ when $i \in N \setminus S$.
Take $i \in S$ (the case $i \in N \setminus S$ is similar and we omit it)

$$\begin{aligned}
 B_i(N_0, C) &= \frac{1}{n!} \sum_{\pi \in \Pi_N} B_i^\pi(N_0, C) \\
 &= \frac{1}{n!} \sum_{\pi \in \Pi_N, \pi_S = \pi'} \left(\sum_{\pi' \in \Pi_S} B_i^{\pi'}(N_0, C) \right) \\
 &= \frac{1}{n!} \frac{n!}{s!} \sum_{\pi' \in \Pi_S} B_i^{\pi'}(N_0, C) = B_i(S_0, C).
 \end{aligned}$$

Then, B satisfies *SEP*.

- *ESEC*. YES. Let (N_0, C) and (N_0, C') be as in the definition of *ESEC*. It is straightforward to see that for all $\pi \in \Pi_N$,

$$B_i^\pi(N_0, C') = \begin{cases} B_i^\pi(N_0, C) + (c'_0 - c_0) & \text{if } i = \pi(1) \\ B_i^\pi(N_0, C) & \text{otherwise.} \end{cases}$$

Now it is easy to conclude that B satisfies *ESEC*.

- *IIT*. NO. See Example 1 and Proposition 5 (a).
- *EC*. NO. It is a trivial consequence of Theorem 2.

About Kar's rule K .

- *CS*. NO. Let (N_0, C) be such that $N = \{1, 2, 3\}$, $c_{01} = 10$, $c_{02} = 18$, $c_{03} = 9$, $c_{12} = 10$, $c_{13} = 10$, and $c_{23} = 9$. Making some computations we obtain that $K(N_0, C) = (8.66, 12.16, 7.16)$ and $v(\{1, 2\}) = 20$.
- *CM*. YES. If c_{ij} increases, $v_C(S \cup i) - v_C(S)$ does not decrease. Then, $K_i(N_0, C) = Sh_i(N, v_C)$ does not decrease.
- *SCM*. NO. See Example 1 (K does not depend on the canonical form) and Proposition 5.
- *PM*. NO. K does not satisfy *CS* and *PM* implies *CS*.
- *CON*. YES because $K(N_0, C) = Sh(N, v_C)$.
- *POS*. NO. See Example 1 with $x = 100$.
- *SEP*. NO. Let (N_0, C) be such that $N = \{1, 2, 3\}$, $c_{01} = 10$, $c_{02} = 100$, $c_{03} = 20$, $c_{12} = 10$, $c_{13} = 100$, and $c_{23} = 40$. Take $S = \{1, 2\}$. Then, $K_1(S_0, C) = -35$ but $K_1(N_0, C) = -15$.
- *ESEC*. YES. Let (N_0, C) and (N_0, C') be as in the definition of *ESEC*. It is easy to see that $v_{C'}(S) = v_C(S) + (c'_0 - c_0)$ for all $S \subset N$. Then,

$$\begin{aligned} K_i(N_0, C) &= Sh_i(N, v_{C'}) = Sh_i(N, v_C) + \frac{c'_0 - c_0}{n} \\ &= K_i(N_0, C) + \frac{c'_0 - c_0}{n}. \end{aligned}$$

- *IIT*. NO. See Example 1 and Proposition 5 (a).
- *EC*. NO. It is a trivial consequence of Theorem 2.

About Dutta-Kar's rule *DK*.

- *CS*. YES. See Dutta and Kar (2002).
- *CM*. YES. See Dutta and Kar (2002).
- *SCM*. NO. See Example 1 (*K* does not depend on the canonical form) and Proposition 5.
- *PM*. NO. Let (N_0, C) be such that $N = \{1, 2, 3\}$ and $c_{01} = 10$, $c_{02} = 100$, $c_{03} = 8$, $c_{12} = 2$, $c_{13} = 5$, and $c_{23} = 4$. Making some computations we obtain that $DK_1(N_0, C) = 8 > 2 = DK_1(\{1, 2\}_0, C)$. Then, *DK* does not satisfy *PM*.
- *CON*. NO. See Example 1.
- *POS*. YES. It is trivial.
- *SEP*. YES. Assume that when we apply Prim's algorithm in (N_0, C) the order induced is $\pi = \{i_1, \dots, i_n\}$. Applying an induction argument over p^q it is easy to prove that if $i_q \in S$ and $x_q = \min \left\{ p^q, c_{i_{q+1}i_{q+1}^*} \right\} = c_{i_{q+1}i_{q+1}^*} < p^q$ then, $i_{q+1} \in S$. This means that "switches costs" among an agent of S and an agent of $N \setminus S$ are not possible.

Now it is easy to see that $DK_i^\pi(N_0, C) = DK_i^{\pi_S}(S_0, C)$ when $i \in S$ and $DK_i^\pi(N_0, C) = DK_i^{\pi_{N \setminus S}}((N \setminus S)_0, C)$ when $i \in N \setminus S$. Since

$$DK_i(N_0, C) = \frac{1}{n!} \sum_{\pi \in \Pi_N} DK_i^\pi(N_0, C)$$

using similar arguments to those used when we proved that *B* satisfies *SEP*, we can conclude that *DK* satisfies *SEP*.

- *ESEC*. NO. Take (N_0, C) and (N_0, C') such that $c_{01} = c_{02} = c_{03} = 10$, $c'_{01} = c'_{02} = c'_{03} = 13$, $c_{12} = c_{13} = c'_{12} = c'_{13} = 10$, and $c_{23} = c'_{23} = 6$. Then, $B(N_0, C) = (10, 8, 8)$ and $B(N_0, C') = (12, 8.5, 8.5)$.

- *IIT*. NO. See Example 1 and Proposition 5 (a).
- *EC*. NO. It is a trivial consequence of Theorem 2.

7.8 Proof of Theorem 2

(a) By Theorem 1 we know that φ satisfies *SEP*, *ESEC*, and *IIT*.

We now prove the uniqueness. Let ψ be a rule satisfying these properties. We apply an induction argument over n . If $n = 1$, $\psi = \varphi$. Assume that the result holds for all $n \leq p - 1$. We prove it for $n = p$.

Since ψ satisfies *IIT*, by Proposition 5 (a), we can restrict to canonical matrices. By simplicity assume that the tree t associated to the canonical matrix C satisfying (A1) is $\{(i-1, i)\}_{i=1}^n$. Let $j \in N$ be such that $c_{(j-1)j} = \max_{i \in N} \{c_{(i-1)i}\}$. We distinguish two cases:

Case 1. $j > 1$. Take $S = \{1, \dots, j-1\}$. By Proposition 3 (a) we know that $t' = \{(i-1, i)\}_{i=1}^{j-1}$ is a *mcst* in (S_0, C) and $t' = \{(0, j)\} \cup \{(i-1, i)\}_{i=j+1}^n$ is a *mcst* in $((N \setminus S)_0, C)$. Then, $m(S_0, C) = \sum_{i=1}^{j-1} c_{(i-1)i}$ and $m((N \setminus S)_0, C) = c_{0j} + \sum_{i=j+1}^n c_{(i-1)i}$.

Since C is a canonical matrix, $c_{0j} = \max_{i \leq j} \{c_{(i-1)i}\} = c_{(j-1)j}$. Then,

$$m(S_0, C) + m((N \setminus S)_0, C) = \sum_{i=1}^n c_{(i-1)i} = m(N_0, C).$$

By *SEP*, $\psi_i(N_0, C) = \psi_i(S_0, C)$ when $i \in S$ and $\psi_i(N_0, C) = \psi_i((N \setminus S)_0, C)$ when $i \notin S$.

We know that $S \neq \emptyset$ and $N \setminus S \neq \emptyset$ because $j > 1$. By induction hypothesis, $\psi(S_0, C) = \varphi(S_0, C)$ and $\psi((N \setminus S)_0, C) = \varphi((N \setminus S)_0, C)$.

Since φ satisfies *SEP* we can conclude that $\psi(N_0, C) = \varphi(N_0, C)$.

Case 2. $j = 1$. Let $k \in N \setminus \{1\}$ be such that $c_{(k-1)k} = \max_{i > 1} \{c_{(i-1)i}\}$. If $c_{(k-1)k} = c_{(j-1)j}$ using similar arguments to those used in Case 1 we can conclude that $\psi(N_0, C) = \varphi(N_0, C)$.

Assume that $c_{(k-1)k} < c_{(j-1)j}$.

Consider now the *cstp* (N_0, C^1) where $c_{il}^1 = c_{il}$ if $0 \notin \{i, l\}$ and $c_{i0}^1 = c_{i0} - x$ where $x = c_{(j-1)j} - c_{(k-1)k} > 0$. Since $c_{i0} = c_{(j-1)j}$ for all $i \in N$ we can apply the property of *ESEC* to problems (N_0, C^1) and (N_0, C) . Then, for all $i \in N$,

$$\begin{aligned}\psi_i(N_0, C) &= \psi_i(N_0, C^1) + \frac{x}{n} \text{ and} \\ \varphi_i(N_0, C) &= \varphi_i(N_0, C^1) + \frac{x}{n}.\end{aligned}$$

It is straightforward to prove that C^1 is a canonical matrix satisfying that $c_{(k-1)k}^1 = \max_{i \in N} \{c_{(i-1)i}^1\}$. Applying Case 1 to C^1 we conclude that $\psi(N_0, C^1) = \varphi(N_0, C^1)$. Then, $\psi(N_0, C) = \varphi(N_0, C)$.

(b) By Theorem 1 we know that φ satisfies *EC*.

We now prove the uniqueness. Let ψ be a rule satisfying *EC*. We prove that $\varphi = \psi$ by induction on n . If $n = 1$ it is trivial to see that $\varphi = \psi$. Assume that the result holds when $n \leq p - 1$ and we prove it when $n = p$.

Given $i, j \in N$, by simplicity we write $\varphi_i = \varphi_i(N_0, C)$, $\psi_i = \psi_i(N_0, C)$, $\varphi_i^{+j} = \varphi_i((N \setminus \{j\})_0, C^{+j})$, and $\psi_i^{+j} = \psi_i((N \setminus \{j\})_0, C^{+j})$.

EC can be written, for ψ , as

$$\psi_i - \psi_i^{+j} = \psi_j - \psi_j^{+i}.$$

Then,

$$\sum_{j \in N \setminus \{i\}} \psi_i - \sum_{j \in N \setminus \{i\}} \psi_i^{+j} = \sum_{j \in N \setminus \{i\}} \psi_j - \sum_{j \in N \setminus \{i\}} \psi_j^{+i}.$$

Since $\sum_{j \in N \setminus \{i\}} \psi_i = (n-1)\psi_i$ and $\sum_{j \in N \setminus \{i\}} \psi_j = m(N_0, C) - \psi_i$,

$$n\psi_i = m(N_0, C) + \sum_{j \in N \setminus \{i\}} \psi_i^{+j} - \sum_{j \in N \setminus \{i\}} \psi_j^{+i}.$$

Since φ also satisfies *EC*,

$$n\varphi_i = m(N_0, C) + \sum_{j \in N \setminus \{i\}} \varphi_i^{+j} - \sum_{j \in N \setminus \{i\}} \varphi_j^{+i}.$$

By induction hypothesis, for all $i, j \in N$, $\psi_i^{+j} = \varphi_i^{+j}$ and $\psi_j^{+i} = \varphi_j^{+i}$. Then, $\varphi_i = \psi_i$ for all $i \in N$.

7.9 Proof of Remark 6

We prove that new rules appear if we remove some of the properties of Corollary 1 or Theorem 2 (a).

IIT in Theorem 2 (a) and *SCM* in Corollary 1. Consider the following subset of permutations:

$$\Pi' = \{\pi \in \Pi_N \mid \pi(i) < \pi(j) \text{ when } c_{0i} < c_{0j}\}.$$

For each *cstp* (N_0, C) and $i \in N$ we define

$$\psi_i^1(N, C) = \frac{1}{|\Pi'|} \sum_{\pi \in \Pi'} (v_{C^*}(Pre(i, \pi) \cup \{i\}) - v_{C^*}(Pre(i, \pi))).$$

It is not difficult to prove that ψ^1 satisfies *PM* (and hence *SEP*) and *ESEC*.

Nevertheless ψ^1 does not satisfy *IIT*. Let (N_0, C) and (N_0, C') be such that $N_0 = \{1, 2, 3\}$, $c_{01} = 2$, $c_{02} = 2$, $c_{03} = 1.5$, $c_{12} = 9$, $c_{13} = 1$, $c_{23} = 1$, $c'_{02} = 3$, and $c'_{ij} = c_{ij}$ if $(i, j) \neq (2, 0)$. Then, $\psi_2^1(N, C') = 1 < 1.5 = \psi_2^1(N, C)$. Since (N_0, C) and (N_0, C') are directly equivalents ($\{(0, 1), (1, 3), (2, 3)\}$ is a *mcst* in both problems) we conclude that ψ^1 does not satisfy *IIT*. Since *SCM* implies *IIT* (Proposition 5 (a)) we conclude that ψ^1 does not satisfy *SCM*.

SEP in Theorem 2 (a) and *PM* in Corollary 1.. Let ψ^2 be the egalitarian rule, *i.e.*

$$\psi_i^2(N, C) = \frac{m(N_0, C)}{n} \text{ for all } i \in N.$$

It is trivial to see that ψ^2 satisfies *SCM* (and hence *IIT*) and *ESEC*.

Let (N_0, C) be such that $N_0 = \{1, 2\}$, $c_{01} = 2$, $c_{02} = 4$, and $c_{12} = 6$. Then, $m(N_0, C) = m(\{1\}_0, C) + m(\{2\}_0, C)$ and $\psi_1^2(\{1\}, C) = 2 \neq 3 = \psi_1^2(N, C)$. Then, ψ^2 does not satisfy *SEP*. Since *PM* implies *SEP* we conclude that ψ^2 does not satisfy *PM*.

ESEC in Theorem 2 (a) and Corollary 1. Given $S \subset \mathcal{N} = \{1, 2, \dots\}$ we denote by π^S the order induced in S by the index of the agents, *i.e.* given $i, j \in S$, $\pi^S(i) < \pi^S(j)$ if and only if $i < j$.

For each $cstp(N_0, C)$ and $i \in N$ we define

$$\psi_i^3(S, C) = v_{C^*}(Pre(i, \pi^S) \cup \{i\}) - v_{C^*}(Pre(i, \pi^S)).$$

It is not difficult to see that ψ^3 satisfies *SCM* (and hence *IIT*) and *PM* (and hence *SEP*).

Nevertheless ψ^2 does not satisfy *ESEC*. Let (N_0, C) be as in Example 1 when $x = 0$ and C' such that $c'_{01} = 12$ and $c'_{02} = 12$. Then, $\psi^3(N, C) = (10, 2)$ whereas $\psi^2(N, C) = (12, 2)$.

7.10 Proof of Proposition 6

(a) Assume that $t = \{(i-1, i)\}_{i=1}^n$ is the tree associated to C satisfying (A1). Take $S = \{i_1, \dots, i_{|S|}\}$ where $i_{p-1} \leq i_p$ for all $p = 2, \dots, |S|$.

For each $p = 1, \dots, |S|$ we consider the following sets

$$S^p = \{i \in N \mid i_{p-1} < i < i_p\}.$$

Moreover,

$$S^{|S|+1} = \begin{cases} \{i \in N \mid i_{|S|} < i \leq n\} & \text{if } i_{|S|} < n \\ \emptyset & \text{otherwise.} \end{cases}$$

In order to simplify the notation we take $i_{|S|+1} = n$.

Notice that, $S^p \subset N \setminus S$ for all $p = 1, \dots, |S| + 1$, $\bigcup_{p=1}^{|S|+1} S^p = N \setminus S$, and $S^p \cap S^q = \emptyset$ for all $p \neq q$. Notice that S^p can be empty.

We know that $v^+(N \setminus S) = m((N \setminus S)_0, C^{+S}) = c((N \setminus S)_0, C^{+S}, t')$ where t' is computed following Prim's algorithm. We now compute t' .

Assume that $i \in S_p$, $j \in S_q$ and $p < q$. Then, $i < i_p \leq i_{q-1} < j$. Since t satisfies (A2),

$$c_{0i}^{+S} = \min_{k \in S_0} \{c_{ik}\} = \min \{c_{i_{p-1}i}, c_{ii_p}\} \leq c_{ii_p}.$$

Moreover, $c_{ij}^{+S} = c_{ij}$. Since t satisfies (A2) we conclude that $c_{ii_p} \leq c_{ij}$ and hence $c_{0i}^{+S} \leq c_{ij}^{+S}$. This means that we can construct t' such that there is no arc linking directly agents of S^p and S^q .

Then, t' can be expressed as $\bigcup_{p=1}^{|S|+1} t'_{S_0^p}$ where $t'_{S_0^p}$ is a *mcst* computed following Prim's algorithm in (S_0^p, C^{+S}) . If $S^p = \emptyset$ we take $t'_{S_0^p} = \emptyset$.

We now apply Prim's algorithm to (S_0^p, C^{+S}) where $p < |S| + 1$. We first select an arc $(0, i)$ such that $c_{0i}^{+S} = \min_{j \in S^p} \{c_{0j}^{+S}\}$. Since t satisfies (A2) we conclude that $c_{0i}^{+S} = \min \{c_{i_{p-1}i}, c_{ii_p}\}$.

Since t satisfies (A2),

$$\min_{j \in S^p} \{c_{0j}^{+S}\} = \min \{c_{i_{p-1}(i_{p-1}+1)}, c_{(i_{p-1})i_p}\} = \min \{c_{0(i_{p-1}+1)}^{+S}, c_{(i_{p-1})0}^{+S}\}.$$

Assume that the arc selected is $(0, i_{p-1} + 1)$ (the case where the arc selected is $(i_p - 1, 0)$ is similar). Then, $c_{0(i_{p-1}+1)}^{+S} = c_{i_{p-1}(i_{p-1}+1)}$, *i.e.* the arc $(0, i_{p-1} + 1)$ in (S_0^p, C^{+S}) is the arc $(i_{p-1}, i_{p-1} + 1)$ in (N_0, C) .

Using arguments similar to those used before we can conclude that the second arc selected is the arc (i, j) such that

$$c_{ij}^{+S} = \min \{c_{(i_{p-1}+1)(i_{p-1}+2)}, c_{(i_{p-1})i_p}\}.$$

If we continue with this procedure we obtain that the cost of the tree $t'_{S_0^p}$ in (S_0^p, C^{+S}) coincide with the cost of the network $t_{S^p \cup \{i_{p-1}, i_p\}} \setminus \{(i_{p'-1}, i_{p'})\}$ in (N_0, C) where $(i_{p'-1}, i_{p'})$ satisfies that

$$c_{i_{p'-1}i_{p'}} = \max_{(i,j) \in t_{S^p \cup \{i_{p-1}, i_p\}}} c_{ij}.$$

Since t satisfies (A2) we conclude that $c_{i_{p'-1}i_{p'}} = c_{i_{p-1}i_p}$. Then,

$$m(S_0^p, C^{+S}) = \sum_{j=i_{p-1}+1}^{i_p} c_{(j-1)j} - c_{i_{p-1}i_p}.$$

Notice that if $S^p = \emptyset$ and we compute $m(S_0^p, C^{+S})$ according with the previous formula we obtain that

$$m((S_p)_0, C^{+S}) = c_{i_{p-1}i_p} - c_{i_{p-1}i_p} = 0.$$

If $p = |S| + 1$ it is trivial to see that

$$m(S_0^{|S|+1}, C^{+S}) = \sum_{j=i_p+1}^n c_{(j-1)j}.$$

Thus,

$$\begin{aligned}
v^+(N \setminus S) &= c((N \setminus S)_0, C^{+S}, t') = \sum_{p=1}^{|S|+1} m((S_p)_0, C^{+S}) \\
&= \sum_{p=1}^{|S|} \left(\sum_{j=i_{p-1}+1}^{i_p} c_{(j-1)j} - c_{i_{p-1}i_p} \right) + \sum_{j=i_p+1}^n c_{(j-1)j} \\
&= \sum_{p=1}^{|S|} \sum_{j=i_{p-1}+1}^{i_p} c_{(j-1)j} + \sum_{j=i_p+1}^n c_{(j-1)j} - \sum_{p=1}^{|S|} c_{i_{p-1}i_p} \\
&= \sum_{j=1}^n c_{(j-1)j} - \sum_{p=1}^{|S|} c_{i_{p-1}i_p}.
\end{aligned}$$

Since t is a *mcst* of (N_0, C) , $\sum_{j=1}^n c_{(j-1)j} = m(N_0, C)$. Moreover, by Proposition 3 (a), $\sum_{p=1}^{|S|} c_{i_{p-1}i_p} = v(S)$. This concludes the proof of (a).

(b) In order to prove that φ^1 satisfies *IIT* it is enough to prove that if (N_0, C) and (N_0, C') are directly equivalent, then $\varphi^1(N_0, C) = \varphi^1(N_0, C')$. Assume that $t = \{(i^0, i)\}_{i=1}^n$ is a *mcst* of (N_0, C) and (N_0, C') satisfying that $c_{i^0i} = c'_{i^0i}$ for all $i = 1, \dots, n$.

We proceed by induction on n . If $n = 1$ it is trivial to see that $\varphi^1(N_0, C) = \varphi^1(N_0, C')$. Assume that the result holds when $n \leq p - 1$. We now prove it when $n = p$.

In order to simplify the notation we take $N^{-i} = N \setminus \{i\}$.

We prove it in several claims.

Claim 1. Given a *cstp* (N_0, C) , $S \subset N$, and $j \in N \setminus S$, $(S_0, C^{+N \setminus S}) = (S_0, (C^{+j})^{+(N^{-j} \setminus S)})$.

Take $i, k \in S$ such that $i \neq 0$ and $k \neq 0$. Then,

$$c_{ik}^{+N \setminus S} = c_{ik} = (c_{ik}^{+j})^{+(N^{-j} \setminus S)}.$$

Given $i \in S$,

$$\begin{aligned}
c_{0i}^{+N \setminus S} &= \min_{k \in (N \setminus S)_0} \{c_{ki}\} \\
&= \min \left\{ \min_{k \in (N^{-j} \setminus S)} \{c_{ki}\}, \min \{c_{0i}, c_{ji}\} \right\} \\
&= \min \left\{ \min_{k \in (N^{-j} \setminus S)} \{c_{ki}^{+j}\}, c_{0i}^{+j} \right\} \\
&= \min_{k \in (N^{-j} \setminus S)_0} \{c_{ki}^{+j}\} \\
&= (c_{0i}^{+j})^{+(N^{-j} \setminus S)}.
\end{aligned}$$

Thus, $C^{+N \setminus S} = (C^{+j})^{+(N^{-j} \setminus S)}$.

Claim 2. Assume that t^* is a *mcst* in (N_0, C) and $j \in N$. Let $g = \{(i_{p-1}, i_p)\}_{p=1}^r$ be the unique path in t^* from $0 (= i_0)$ to $j (= i_r)$. Assume that $c_{i_{q-1}i_q} = \max_{p=1, \dots, r} \{c_{i_{p-1}i_p}\}$. Then, $t^* \setminus \{(i_{q-1}, i_q)\}$ is a *mcst* in (N_0^{-j}, C^{+j}) .

We denote $t_q^* = t^* \setminus \{(i_{q-1}, i_q)\}$. Then, t_q^* can be identified with a tree in (N_0^{-j}, C^{+j}) simply by changing the arc (k, j) , when $(k, j) \in t_q^*$, by $(k, 0)$.

Suppose that t_q^* is not a *mcst* in (N_0^{-j}, C^{+j}) . Then, it exists a tree t' in (N_0^{-j}, C^{+j}) such that

$$c(N_0^{-j}, C^{+j}, t') < c(N_0^{-j}, C^{+j}, t_q^*).$$

Let S_j denote the set of agents of N^{-j} who are connected to the source in t' through agent j . We now define S_j formally. For each $i \in N^{-j}$, let $\{(0, l_1), (l_1, l_2), \dots, (l_{p-1}, i)\}$ be the unique path in t' from the source to i . Then,

$$S_j = \{i \in N^{-j} \mid c_{0l_1}^{+j} = c_{jl_1} \text{ and } (j, l_1) \in t^*\}.$$

Notice that there is no arc $(i, k) \in t'$ such that $i \in S_j$, $k \in N^{-j} \setminus S_j$. Otherwise, t' have a cycle and could not be a tree in (N_0^{-j}, C^{+j}) . We can find $(i_{r-1}, i_r) \in g \subset t^*$ such that $i_{r-1} \in (N^{-j} \setminus S_j) \cup \{0\}$ and $i_r \in S_j \cup \{j\}$.

If we change the arcs $(0, l) \in t'$ such that $c_{0l_1}^{+j} = c_{jl_1}$ and $(j, l_1) \in t^*$ by the arc (j, l_1) , $t' \cup \{(i_{r-1}, i_r)\}$ could be identified with a tree in (N_0, C) .

Since

$$\begin{aligned} c(N_0, C, t' \cup \{(i_{r-1}, i_r)\}) &= c(N_0^{-j}, C^{+j}, t') + c_{i_{r-1}i_r}, \\ c(N_0, C, t^*) &= c(N_0^{-j}, C^{+j}, t_q^*) + c_{i_{q-1}i_q}, \text{ and} \\ c_{i_{r-1}i_r} &\leq c_{i_{q-1}i_q} \end{aligned}$$

we conclude that

$$c(N_0, C, t' \cup \{(i_{r-1}, i_r)\}) < c(N_0, C, t^*),$$

which is a contradiction because t^* is a *mcst* of (N_0, C) .

Claim 3. Take $j \in N$. For all $i \in N^{-j}$

$$\varphi_i^1(N^{-j}, C^{+j}) = \varphi_i^1(N^{-j}, C'^{+j}).$$

Let $g = \{(i_{p-1}, i_p)\}_{p=1}^r$ be the unique path in t from $0 (= i_0)$ to $j (= i_r)$. Take $c_{k^0k} = \max_{p=1, \dots, r} \{c_{i_{p-1}i_p}\}$. Since t is a *mcst* in (N_0, C) and (N_0, C') we conclude, by Claim 2, that $t \setminus \{(k^0, k)\}$ is a *mcst* of (N_0^{-j}, C^{+j}) and (N_0^{-j}, C'^{+j}) satisfying that $c_{i^0i} = c'_{i^0i}$ for all $i \in N \setminus \{k\}$. Then, (N_0^{-j}, C^{+j}) and (N_0^{-j}, C'^{+j}) are directly equivalent.

By induction hypothesis, $\varphi_i^1(N^{-j}, C^{+j}) = \varphi_i^1(N^{-j}, C'^{+j})$.

Claim 4. Given a *cstp* (N_0, C) and $i, j \in N$ such that $i \neq j$,

$$\varphi_i^1(N^{-j}, C^{+j}) = Sh_i(N^{-j}, v^+).$$

For all $S \subset N^{-j}$,

$$\begin{aligned} v_C^+(S) &= m(S_0, C^{+N \setminus S}) \text{ and} \\ v_{C^{+j}}^+(S) &= m\left(S_0, (C^{+j})^{+((N \setminus j) \setminus S)}\right). \end{aligned}$$

By Claim 1, $v_C^+(S) = v_{C^{+j}}^+(S)$. Then,

$$\varphi_i^1(N^{-j}, C^{+j}) = Sh_i(N^{-j}, v_{C^{+j}}^+) = Sh_i(N^{-j}, v_C^+).$$

Claim 5. For all $i, j \in N$, $i \neq j$,

$$Sh_i(N^{-j}, v_C^+) = Sh_i(N^{-j}, v_{C'}^+).$$

By Claim 4, $Sh_i(N^{-j}, v_C^+) = \varphi_i^1(N^{-j}, C^{+j})$ and $Sh_i(N^{-j}, v_{C'}^+) = \varphi_i^1(N^{-j}, C'^{+j})$. By Claim 3, $\varphi_i^1(N^{-j}, C^{+j}) = \varphi_i^1(N^{-j}, C'^{+j})$. Thus, Claim 5 holds.

Claim 6. For all $i \in N$, $\varphi_i^1(N_0, C) = \varphi_i^1(N_0, C')$.

In Pérez-Castrillo and Wettstein (2001) appears the following expression of the Shapley value:

$$Sh_i(N, w) = \frac{1}{n} \left[w(N) - w(N^{-i}) + \sum_{j \in N^{-i}} Sh_i(N^{-j}, w) \right].$$

Since $w(N^{-i}) = \sum_{j \in N^{-i}} Sh_j(N^{-i}, w)$ we conclude that

$$Sh_i(N, w) = \frac{1}{n} \left[w(N) - \sum_{j \in N^{-i}} Sh_j(N^{-i}, w) + \sum_{j \in N^{-i}} Sh_i(N^{-j}, w) \right].$$

Since $\varphi^1(N_0, C) = Sh(N, v_C^+)$ and $\varphi^1(N_0, C') = Sh(N, v_{C'}^+)$,

$$\begin{aligned} \varphi_i^1(N_0, C) &= \frac{1}{n} \left[v_C^+(N) - \sum_{j \in N^{-i}} Sh_j(N^{-i}, v_C^+) + \sum_{j \in N^{-i}} Sh_i(N^{-j}, v_C^+) \right] \text{ and} \\ \varphi_i^1(N_0, C') &= \frac{1}{n} \left[v_{C'}^+(N) - \sum_{j \in N^{-i}} Sh_j(N^{-i}, v_{C'}^+) + \sum_{j \in N^{-i}} Sh_i(N^{-j}, v_{C'}^+) \right]. \end{aligned}$$

Since $v_C^+(N) = v_{C'}^+(N) = m(N_0, C) = m(N_0, C')$ and Claim 5 we conclude that $\varphi_i^1(N_0, C) = \varphi_i^1(N_0, C')$.

(c) We must prove that for all $i, j \in N$, $i \neq j$,

$$\varphi_i^1(N_0, C) - \varphi_i^1(N_0^{-j}, C^{+j}) = \varphi_j^1(N_0, C) - \varphi_j^1(N_0^{-i}, C^{+i}).$$

Given a *TU* game (N, w) Myerson (1980) proved that the Shapley value satisfies that for all $i, j \in N$, $i \neq j$,

$$Sh_i(N, w) - Sh_i(N^{-j}, w) = Sh_j(N, w) - Sh_j(N^{-i}, w).$$

We know that $\varphi^1(N_0, C) = Sh(N, v^+)$. By Claim 4 of part (b), $\varphi_i^1((N^{-j})_0, C^{+j}) = Sh_i(N^{-j}, v^+)$ and $\varphi_j^1((N^{-i})_0, C^{+i}) = Sh_j(N^{-i}, v^+)$.

Applying Myerson's result to the *TU* game (N, v^+) we obtain that φ^1 satisfies *EC*.

7.11 Proof of Theorem 3

We prove several claims:

Claim 1. $\varphi^1 = \varphi$.

By definition, $\varphi(N_0, C) = K(N_0, C^*) = Sh(N, v_{C^*})$. By Proposition 6 (b) and Proposition 5 (a) we conclude that $\varphi^1(N_0, C) = \varphi^1(N_0, C^*) = Sh(N, v_{C^*}^+)$.

By Proposition 6 (a), $v_{C^*}(S) + v_{C^*}^+(N \setminus S) = m(N_0, C)$ for all $S \subset N$. Since $v_{C^*}(N) = v_{C^*}^+(N) = m(N_0, C^*)$ and the Shapley value of a *TU* game (N, w) can be expressed for all $i \in N$ as

$$Sh_i(N, w) = \frac{1}{n!} \sum_{\pi \in \Pi_N} (w(Pre(i, \pi) \cup \{i\}) - w(Pre(i, \pi))),$$

it is easy to conclude that $Sh(N, v_{C^*}) = Sh(N, v_{C^*}^+)$. Thus, $\varphi(N_0, C) = K(N_0, C)$.

Claim 2. $\varphi = \varphi^3$.

We must prove that for all $i \in N$, $\varphi_i(N_0, C) = \frac{1}{n!} \sum_{\pi \in \Pi_N} c_{i^\pi}^*$. Since φ satisfies *IIT* it is enough to prove that if C is a canonical matrix then, $\varphi_i(N_0, C) = \frac{1}{n!} \sum_{\pi \in \Pi_N} c_{i^\pi}$.

We proceed in several claims.

Claim 2a. Let (N_0, C) be a *cstp* where C is a canonical matrix and $t = \{(i_{p-1}, i_p)\}_{p=1}^n$ is the tree given by (A1). Assume that $c_{i_{p-1}i_p} \neq c_{i_{q-1}i_q}$ for all p, q such that $p \neq q$. Given $\pi \in \Pi_N$ we can find a one-to-one application $f_\pi : N \rightarrow t$ such that $f_\pi(i) = (i_{p-1}, i_p)$ satisfies $v(Pre(i, \pi) \cup \{i\}) - v(Pre(i, \pi)) = c_{i_{p-1}i_p} = c_{i^\pi}$.

By simplicity we take $\pi = (1, 2, \dots, n)$. Assume that $1 = i_q$. By Proposition 3 (b) we know that

$$v(Pre(1, \pi) \cup \{1\}) - v(Pre(1, \pi)) = v(1) = c_{i_0 i_q}.$$

Since C is a canonical matrix we can find $(i_{p_1-1}, i_{p_1}) \in t$ such that $p_1 \leq q$ and $c_{i_0 i_q} = c_{i_{p_1-1} i_{p_1}}$. Moreover $1^\pi = 0$ and $c_{01} = c_{i_0 i_q} = c_{i_{p_1-1} i_{p_1}}$. We define $f_\pi(1) = (i_{p_1-1}, i_{p_1})$.

Assume that we have defined $f_\pi(j)$ for all $j \leq i-1$ satisfying that $f_\pi(j) = (i_{p_{j-1}}, i_{p_j}) \in t$ for all $j \leq i-1$ and $f_\pi(j) \neq f_\pi(k)$ for all $j \leq i-1, k \leq i-1, j \neq k$.

We now define $f_\pi(i)$. Assume that $i = i_q$ and $Pre(i, \pi) = \{i_{q_1}, i_{q_2}, \dots, i_{q_{i-1}}\}$ where $q_{l-1} \leq q_l$ for all $l = 2, \dots, i-1$. Notice that we order the agents of $Pre(i, \pi)$ according with t . Two cases can occur:

Firstly, $q < q_{i-1}$. By Proposition 3 (b),

$$v(Pre(i, \pi) \cup \{i\}) - v(Pre(i, \pi)) = \min \left\{ c_{i_{q_{k-1}}i_q}, c_{i_qi_{q_k}} \right\}$$

where $q_{k-1} < q < q_k$ and $k \leq i-1$.

Since t satisfies (A2) and $c_{i_{p-1}i_p} \neq c_{i_{q-1}i_q}$ for all p, q such that $p \neq q$ we conclude that $c_{i_{q_{k-1}}i_q} \neq c_{i_qi_{q_k}}$. Assume that $\min \left\{ c_{i_{q_{k-1}}i_q}, c_{i_qi_{q_k}} \right\} = c_{i_qi_{q_k}}$ (the other case is similar).

By Proposition 3 (b), for all $j \leq i$

$$c_{i_qi_{q_k}} \neq v(Pre(j, \pi) \cup \{j\}) - v(Pre(j, \pi)).$$

Then, we can find $(i_{p_{i-1}}, i_{p_i}) \in t \setminus \left(\bigcup_{j=1}^{i-1} (i_{p_{j-1}}, i_{p_j}) \right)$ such that $q \leq p_i - 1 < p_i \leq q_k$ and $c_{i_{p_{i-1}}i_{p_i}} = c_{i_qi_{q_k}}$. Moreover, $c_{i^{\pi i}} = \min_{j \in Pre(i, \pi)} \{c_{ji}\} = c_{i_qi_{q_k}}$.

We now define $f_\pi(i) = (i_{p_{i-1}}, i_{p_i})$.

Secondly, $q > q_{i-1}$. By Proposition 3 (b),

$$v(Pre(i, \pi) \cup \{i\}) - v(Pre(i, \pi)) = c_{i_{q_{i-1}}i_q}.$$

By Proposition 3 (b) we can find $(i_{p_{i-1}}, i_{p_i}) \in t \setminus \left(\bigcup_{j=1}^{i-1} (i_{p_{j-1}}, i_{p_j}) \right)$ such that $q_{i-1} \leq p_i - 1 < p_i \leq q$ and $c_{i_{p_{i-1}}i_{p_i}} = c_{i_{q_{i-1}}i_q}$. Moreover, $c_{i^{\pi i}} = \min_{j \in Pre(i, \pi)} \{c_{ji}\} = c_{i_{q_{i-1}}i_q}$.

We now define $f_\pi(i) = (i_{p_{i-1}}, i_{p_i})$.

Claim 2b. Let (N_0, C) be a *cstp* where C is a canonical matrix and $t = \{(i_{p-1}, i_p)\}_{p=1}^n$ is the tree given by (A1). Assume that $c_{i_{p-1}i_p} = c_{i_{q-1}i_q}$ for some p, q such that $p \neq q$. Given $\pi \in \Pi_N$ we can find a one-to-one application $f_\pi : N \rightarrow t$ such that $f_\pi(i) = (i_{p-1}, i_p)$ satisfies $v(Pre(i, \pi) \cup \{i\}) - v(Pre(i, \pi)) = c_{i_{p-1}i_p} = c_{i^{\pi i}}$.

Let C^ε be a canonical matrix obtained from C modifying slightly the cost of the arcs $c_{i_{p-1}i_p}$ such that there exist $q \neq p$ with $c_{i_{p-1}i_p} = c_{i_{q-1}i_q}$. We modify these cost in such a way that if $c_{i_{p-1}i_p} < c_{i_{q-1}i_q}$ then, $c_{i_{p-1}i_p}^\varepsilon < c_{i_{q-1}i_q}^\varepsilon$ and if $c_{i_{p-1}i_p} = c_{i_{q-1}i_q}$ then $c_{i_{p-1}i_p}^\varepsilon \neq c_{i_{q-1}i_q}^\varepsilon$.

By Claim 2a, given the *cstp* (N_0, C^ε) and $\pi \in \Pi_N$ we can find a one-to-one application $f_\pi^\varepsilon = N \rightarrow t$ such that $f_\pi^\varepsilon(i) = (i_{p-1}, i_p)$ satisfy $v^\varepsilon(Pre(i, \pi) \cup \{i\}) - v^\varepsilon(Pre(i, \pi)) = c_{i_{p-1}i_p}^\varepsilon = c_{ii^\pi}^\varepsilon$.

Given the *cstp* (N_0, C) and $\pi \in \Pi_N$ we define $f_\pi = f_\pi^\varepsilon$. Of course, f_π is a one-to-one application. Since v is a continuous function of C we conclude that

$$v(Pre(i, \pi) \cup \{i\}) - v(Pre(i, \pi)) = c_{i_{p-1}i_p} = c_{ii^\pi}.$$

Claim 2c. If C is a canonical matrix, $\varphi_i(N_0, C) = \frac{1}{n!} \sum_{\pi \in \Pi_N} c_{ii^\pi}$ for all $i \in N$.

Since $\varphi(N_0, C) = Sh(N, v)$, for all $i \in N$,

$$\begin{aligned} \varphi_i(N_0, C) &= \frac{1}{n!} \sum_{\pi \in \Pi_N} (v(Pre(i, \pi) \cup \{i\}) - v(Pre(i, \pi))) \\ &= \frac{1}{n!} \sum_{\pi \in \Pi_N} c_{ii^\pi}. \end{aligned}$$

Claim 3. $\varphi = \varphi^2$.

Before proving Claim 3 we need an additional result. Let (N_0, C) be a *cstp* where C is a canonical matrix and $t = \{(i_{p-1}, i_p)\}_{p=1}^n$ is the tree given by (A1). For each $i \in N$ and $p = 1, \dots, n$ we define

$$\Pi_i^p = \{\pi \in \Pi_N \mid f_\pi(i) = (i_{p-1}, i_p)\} \text{ and } \mu_i^p = \frac{|\Pi_i^p|}{n!}.$$

Claim 3a. Let (N_0, C) be a *cstp*. Given a *mcst* $t^0 = \{(j^0, j)\}_{j \in N}$, for all $i \in N$

$$\varphi_i(N_0, C) = \sum_{p=1}^n \mu_i^p c_{j^0, j}.$$

where for all $i \in N$, $\sum_{j=1}^n \mu_i^j = 1$ and for all $j \in N$, $\sum_{i=1}^n \mu_i^j = 1$.

Let $t = \{(i_{p-1}, i_p)\}_{p=1}^n$ be the *mcst* associated to C^* given by the algorithm when we start with the tree t^0 .

We define the one-to-one application $g : t^0 \rightarrow t$ such that $g(j^0, j) = (i_{p-1}, i_p)$ where $j = i_p$.

By claims 2a, 2b, and 2c, for all $i \in N$, $\varphi_i(N_0, C) = \sum_{p=1}^n \mu_i^p c_{i_{p-1}i_p}^*$; $\sum_{p=1}^n \mu_i^p = 1$ for all $i \in N$; and $\sum_{i=1}^n \mu_i^p = 1$ for all $p = 1, \dots, n$.

We proceed by induction on the number of agents. If $n = 1$ then $\varphi = \varphi^2$. Assume that the result holds when $n \leq p - 1$. We now prove for $n = p$.

Let $t = \{(i^0, i)\}_{i \in N}$ be a *mcst* in (N_0, C) . We denote by $\varphi^2(N, t)$ the rule φ^2 computed when the set of agents is N and the tree starting the process is t . We will prove that $\varphi^2(N, t) = \varphi(N_0, C)$. Notice that this imply that φ^2 is well defined because φ is independent of t .

Take $j \in N$ such that $c_{j^0j} = \max_{i \in N} \{c_{i^0i}\}$. We assume that $c_{j^0j} > c_{i^0i}$ for all $i \in N \setminus \{j\}$. Otherwise we can achieve the same conclusion using similar ideas but in a more elaborate way.

We define $N' = \{i \in N \mid i^0 = 0\}$. We consider three cases:

Case 1. $|N'| \geq 2$. For any $i \in N'$ let F^i be the set of agents $j \in N$ such that $(i, 0)$ is in the unique path from j to 0. Then, $\{F^i\}_{i \in N'}$ is a partition of N satisfying that $\sum_{i \in N'} m((F^i)_0, C) = m(N_0, C)$ and t_{F^i} is a tree in (F^i, C) .

Since φ satisfies *SEP*, for all $i \in N'$ and $k \in F^i$, $\varphi_k(N_0, C) = \varphi_k(F_0^i, C)$.

Because of the process defining φ^2 it is straightforward to prove that for all $i \in N'$ and $k \in F^i$, $\varphi_k^2(N, t) = \varphi_k^2(F^i, t_{F^i})$.

Since $|N'| \geq 2$ we can apply the induction hypothesis to each F^i . Then, for all $i \in N'$ and $k \in F^i$, $\varphi_k^2(F_0^i, t_{F^i}) = \varphi_k(F_0^i, C)$.

This means that φ^2 is well defined and $\varphi^2(N_0, C) = \varphi(N_0, C)$.

Case 2. $|N'| = 1$ and $j^0 \neq 0$. Let F be the set of agents $i \in N$ such that the arc (j^0, j) is in the unique path from i to 0 and $B = N \setminus F$. Notice that $F \neq \emptyset$ and $B \neq \emptyset$ because $j \in F$ and $j^0 \in B$.

Next statements are a trivial consequence of the process defining φ^2 . If $a^s(i) = (j^0, j)$ for some step s and $i \in N$ then the only arc in $A^{s-1} = \{(j^0, j)\}$.

Since in t_B there is exactly $|B|$ arcs, $\sum_{s=1}^{\gamma} p^s = 1$, and the previous statements we conclude that the arcs of t_B are paid by agents in B and the arcs

in $t \setminus t_B$ are paid by agents in F .

Now it is easy to conclude that $\varphi_i^2(N, t) = \varphi_i^2(B, t_B)$ for all $i \in B$. Moreover, if we take j^0 as the source in $(F, t \setminus t_B)$, $\varphi_i^2(N, t) = \varphi_i^2(F, t \setminus t_B)$ for all $i \in F$.

We can compute the canonical form C^* associated to (N_0, C) starting with the tree t . Then, we obtain that the tree $\{(i_{q-1}, i_q)\}_{q=1}^n$ associated to C^* satisfies that it exists $i_{|B|}$ such that $B = \{i_1, \dots, i_{|B|}\}$, $F = \{i_{|B|+1}, \dots, i_n\}$, $j^0 = i_{|B|}$, and $j = i_{|B|+1}$.

It is straightforward to prove that $m(N_0, C^*) = m(B_0, C^*) + m(F_0, C^*)$, t_B is a *mcst* in (B_0, C^*) and $t \setminus t_B$ is a *mcst* in (F_0, C^*) .

Since φ satisfies *SEP*, $\varphi_i(N_0, C) = \varphi_i(B_0, C^*)$ for all $i \in B$ and $\varphi_i(N_0, C) = \varphi_i(F_0, C^*)$ for all $i \in F$.

By induction hypothesis we conclude that $\varphi_i^2(B, t_B) = \varphi_i(B_0, C^*)$ for all $i \in B$ and $\varphi_i^2(F, t \setminus t_B) = \varphi_i(F_0, C^*)$ for all $i \in F$.

Then, φ^2 is well defined and $\varphi^2(N_0, C) = \varphi(N_0, C)$.

Case 3. $|N'| = 1$ and $j^0 = 0$. Take $k \in N \setminus \{j\}$ such that $c_{k^0 k} = \max_{i \in N \setminus \{j\}} \{c_{i^0 i}\}$. We assume that $c_{k^0 k} > c_{i^0 i}$ for all $i \in N \setminus \{j, k\}$. Otherwise we can achieve the same conclusion using similar ideas but in a more elaborate way.

Let F be the set of agents $i \in N$ such that the arc (k^0, k) is in the unique path from i to 0 and $B = N \setminus F$. Notice that $F \neq \emptyset$ and $B \neq \emptyset$ because $k \in F$ and $k^0 \in B$.

We assume that $|B| \geq |F|$. The case $|B| < |F|$ can be proved in a similar way.

Next statements are a trivial consequence of the process defining φ^2 . If $a^s(i) = (0, j)$ for some Step s and $i \in N$ then $A^{s-1} = \{(0, j)\}$. If $a^s(i) = (k^0, k)$ for some Step s and $i \in N$ then $A^{s-1} = \{(0, j), (k^0, k)\}$. Moreover, the cost of the arcs in t_F are paid only by agents in F and the cost of arcs in $t_B \setminus \{(0, j)\}$ are paid only by agents in B .

Thus, we can find $\alpha, \alpha' < \gamma$ such that: In Step α , for all $(i, i') \in t_F$, $\varrho(i, i') = 1$, $\varrho(k^0, k) = \varrho(0, j) = 0$, and $\sum_{s=1}^{\alpha} p^s = \frac{|F|-1}{|F|}$. In Stage α' for all $(i, i') \in t_B$, $\varrho(i, i') = 1$, $\varrho(0, j) = 0$, and $\sum_{s=1}^{\alpha'} p^s = \frac{|B|-1}{|B|}$. Notice that $\alpha = \alpha'$ if and only if $|B| = |F|$.

We can compute the canonical form C^* starting with the tree t . Then we

obtain that the tree $\{(i_{q-1}, i_q)\}_{q=1}^n$ associated to C^* satisfies that it exists $i_{|B|}$ such that $B = \{i_1, \dots, i_{|B|}\}$, $F = \{i_{|B|+1}, \dots, i_n\}$, $k^0 = i_{|B|}$, and $k = i_{|B|+1}$.

By Claim 3a, for all $i \in N$,

$$\varphi_i(N_0, C) = \sum_{q=1}^n \mu_i^q c_{q^0 q}$$

where $\mu_i^q = \frac{|\Pi_i^q|}{n!}$ and

$$\Pi_i^q = \{\pi \in \Pi_N \mid v(\text{Pre}(i, \pi) \cup \{i\}) - v(\text{Pre}(i, \pi)) = c_{q^0 q}\}.$$

Since φ^2 can be expressed as $\sum_{q=1}^n \lambda_i^q c_{q^0 q}$ we only need to prove that for all $(q^0, q) \in t$, $\lambda_i^q = \mu_i^q$ for all $i \in N$.

Consider the problem (F_0, C^*) . For each $i, q \in F$ we denote by $\mu_i^q(F)$ the corresponding function associated to (F_0, C^*) . It is straightforward to prove that for all $q \in F \setminus \{k\}$ and $i \in F$, $\mu_i^q(F) = \mu_i^q$. Moreover, for all $q \in F \setminus \{k\}$ and $i \in B$, $\mu_i^q = 0$.

We know that $t_F \cup \{(0, k)\}$ is a *mcst* in (F_0, C^*) and $c_{0k}^* = c_{0j} > c_{k^0 k}$. For each $i, q \in F_0$ we denote by $\lambda_i^q(F)$ the corresponding function associated to $\varphi^2(F, t_F \cup \{(0, k)\})$. We saw before that when we compute $\varphi^2(N, t)$ the cost of the arcs of t_F are paid only by agents in F . Then, for all $q \in F \setminus \{k\}$ and $i \in F$, $\lambda_i^q(F) = \lambda_i^q$. Moreover, for all $q \in F \setminus \{k\}$ and $i \in B$, $\lambda_i^q = 0$.

Take $q \in F \setminus \{k\}$ and $i \in F$. By induction hypothesis we know that $\mu_i^q(F) = \lambda_i^q(F)$ and hence, $\mu_i^q = \lambda_i^q$. Then, for all $q \in F \setminus \{k\}$ and $i \in N$, $\mu_i^q = \lambda_i^q$.

Using similar arguments to those used with F we can conclude that for all $q \in B \setminus \{j\}$ and $i \in N$, $\mu_i^q = \lambda_i^q$.

We only need to prove that $\mu_i^q = \lambda_i^q$ when $q \in \{j, k\}$ and $i \in N$. It is straightforward to prove that for all $i \in N$, $\mu_i^j = \lambda_i^j = \frac{1}{n}$.

We now prove that for all $i \in N$, $\mu_i^k = \lambda_i^k$.

Take $i \in F$. Π_i^k corresponds to the permutations satisfying that, in the order given by π , the first agent belongs to B and i is the first agent of F . It is straightforward to prove that $\mu_i^k = \frac{|B|}{n|F|}$. Applying a similar argument to B we obtain that $\mu_i^k = \frac{|F|}{n|B|}$ if $i \in B$.

Take $i \in F$. From Step α to Step α' we have that $a^s(i) = (k^0, k)$ for all $i \in F$ whereas $a^s(i) \neq (k^0, k)$ for all $i \in B$. This means that from Step α to Step α' each agent $i \in F$ pays $\left(\frac{|B|-1}{|B|} - \frac{|F|-1}{|F|}\right) c_{k^0k}$. Then,

$$\varrho^{\alpha'}(k, k^0) = \left(1 - |F| \left(\frac{|B|-1}{|B|} - \frac{|F|-1}{|F|}\right)\right)$$

In Step $\alpha' + 1$, $a^{\alpha'+1}(i) = (k^0, k)$ for all $i \in N$. Then, each agent $i \in N$ pays

$$\frac{1}{n} \left(1 - |F| \left(\frac{|B|-1}{|B|} - \frac{|F|-1}{|F|}\right)\right)$$

and $\varrho^{\alpha'+1}(k^0, k) = 1$, which means that the cost of this arc is already paid. Then,

$$\begin{aligned} \lambda_i^k &= \frac{1}{n} \left(1 - |F| \left(\frac{|B|-1}{|B|} - \frac{|F|-1}{|F|}\right)\right) \text{ if } i \in B \text{ and} \\ \lambda_i^k &= \left(\frac{|B|-1}{|B|} - \frac{|F|-1}{|F|}\right) + \frac{1}{n} \left(1 - |F| \left(\frac{|B|-1}{|B|} - \frac{|F|-1}{|F|}\right)\right) \text{ if } i \in F. \end{aligned}$$

Making some computations we obtain that $\lambda_i^k = \mu_i^k$ for all $i \in N$. Then, φ^2 is well defined and $\varphi^2(N_0, C) = \varphi(N_0, C)$.

Claim 4. $\varphi = P$.

Branzei *et al* (2003) characterizes P as the only rule satisfying efficiency (EF), upper bound contributions (UBC), equal treatment (ET), and cone-wise positive linearity (CPL). We prove that φ satisfies these four properties.

EF means that $\sum_{i \in N} \varphi_i(N_0, C) = m(N_0, C)$ for all $cstp(N_0, C)$. Of course, φ satisfies it.

Given a $cstp(N_0, C)$, i and j are (C, S) -connected if there exists a path g from i to j satisfying that for all $(k, l) \in g$, $\{k, l\} \subset S_0 \subset N_0$ and $c_{kl} = 0$. $S \subset N$ is a C -component if two conditions hold. First, for all $i, j \in S$, i and j are (C, S) -connected. Second, if $S \not\subset T$ there exists $i, j \in T$ such that i and j are not (C, T) -connected.

UBC means that for all C -component S , $\sum_{i \in S} \varphi_i(N_0, C) \leq \min_{j \in S} \{c_{0j}\}$.

Let $P = \{S_1, \dots, S_p\}$ be the partition induced in N by C -component coalitions. φ satisfies ET if for all $S_q \in P$ and all $i, j \in S_q$, $\varphi_i(N_0, C) = \varphi_j(N_0, C)$.

Let C^* be the canonical form associated to C . Because of the definition of the algorithm it is trivial to see that if S is a C - component and $t = \{(i_{p-1}, i_p)\}_{p=1}^n$ is the tree satisfying (A1), there exist $i_q \in N$ such that $S = \{i_q, i_{q+1}, \dots, i_{q+|S|-1}\}$, $c_{0i_q}^* \leq \min_{j \in S} \{c_{0j}\}$, and $c_{i_{p-1}i_p}^* = 0$ for all $p = q+1, \dots, q+|S|-1$.

Take $\pi \in \Pi_N$ and $i \in S$, by Proposition 3 (b), $v_{C^*}(Pre(i, \pi) \cup \{i\}) - v_{C^*}(Pre(i, \pi)) \leq c_{0i_q}^*$ if $Pre(i, \pi) \cap S = \emptyset$ and $v_{C^*}(Pre(i, \pi) \cup \{i\}) - v_{C^*}(Pre(i, \pi)) = 0$ otherwise.

We now conclude that φ satisfies *UBC* and *ET* because $\varphi(N_0, C) = Sh(N, v_{C^*})$.

CPL. Let (N_0, C) and (N_0, C') be two *cstp* such that given $i, j, k, l \in N_0$, $c_{ij} \leq c_{kl}$ if and only if $c'_{ij} \leq c'_{kl}$. Take $x, x' \geq 0$. φ satisfies *CPL* if $\varphi(N_0, xC + x'C') = x\varphi(N_0, C) + x'\varphi(N_0, C')$ where xC is the cost matrix defined by $(xC)_{ij} = xc_{ij}$ for all $i, j \in N_0$.

Let t^0 be a *mcst* in (N_0, C) . Then, t^0 is a *mcst* in (N_0, C') and $(N_0, xC + x'C')$. Assume that $t = \{(i_{p-1}, i_p)\}_{p=1}^n$ is the tree satisfying (A1) given by the algorithm when we apply it to C . It is trivial to see that t is also the tree satisfying (A1) given by the algorithm when we apply it to C' and $xC + x'C'$.

Let us denote by C^* , C'^* , and $(xC + x'C')^*$ the canonical matrices computed applying the algorithm when we start in t^0 . Again, it is straightforward to prove that for all $p = 1, \dots, n$

$$(xC + x'C')^*_{i_{p-1}i_p} = x c_{i_{p-1}i_p}^* + x' c'_{i_{p-1}i_p}.$$

By Proposition 3 (a) we conclude that for all $S \subset N$,

$$v_{(xC+x'C')^*}(S) = x v_{C^*}(S) + x' v_{C'^*}(S).$$

Since the Shapley value is additive on the characteristic function we conclude that φ satisfies *CPL*.

8 References

Bergantiños G. and Lorenzo L. (2003). "A non-cooperative approach to the cost spanning tree problem." *Mathematical Methods of Operations Research*, forthcoming.

- Bergantiños G. and Vidal-Puga J. (2004). "Realizing efficient outcomes in cost spanning tree problems". Mimeo.
- Bird C.G. (1976). On cost allocation for a spanning tree: A game theoretic approach. *Networks* 6: 335-350.
- Branzei R., Moretti S., Norde H., and Tijs S. (2003). The P-value for cost sharing in minimum cost spanning tree situations. Mimeo, Tilburg University.
- Dutta B. and Kar A. (2002). Cost monotonicity, consistency and minimum cost spanning tree games. *Games and Economic Behavior*. Forthcoming.
- Feltkamp V., Tijs S., and Muto S. (1994a). On the irreducible core and the equal remaining obligation rule of minimum cost extension problems. Mimeo, Tilburg University.
- Feltkamp V., Tijs S., and Muto S. (1994b). Minimum cost spanning extension problems: the proportional rule and the decentralized rule. Mimeo, Tilburg University.
- Granot D., and Huberman G. (1981). Minimum cost spanning tree games. *Mathematical Programming* 21: 1-18.
- Granot D., and Huberman G. (1984). On the core and nucleolus of the minimum cost spanning tree games. *Mathematical Programming* 29: 323-347.
- Granot D., Maschler M. (1998). Spanning network games. *International Journal of Game Theory* 27: 467-500.
- Kar A (2002). Axiomatization of the Shapley value on minimum cost spanning tree games. *Games and Economic Behavior* 38: 265-277.
- Kruskal J. (1956). On the shortest spanning subtree of a graph and the traveling salesman problem. *Proceedings of the American Mathematical Society* 7: 48-50.
- Megiddo N. (1978). Computational complexity and the game theory approach to cost allocation for a tree. *Mathematics of Operations Research* 3: 189-196.
- Myerson R.B. (1980). Conference structures and fair allocation rules. *International Journal of Game Theory* 9: 169-182.
- Pérez-Castrillo D. and Wettstein D. (2001). Bidding for the surplus: A non-cooperative approach to the Shapley value. *Journal of Economic Theory* 100: 274-294.
- Prim R.C. (1957). Shortest connection networks and some generalizations. *Bell Systems Technology Journal* 36: 1389-1401.

Shapley L.S. (1953). A value for n-person games. In: Kuhn H.W., Tucker A.W. (eds) Contributions to the Theory of Games II, Princeton University Press, Princeton NJ: 307-317.

Sharkey W.W. (1995). Networks models in economics. In Handbooks of operation research and management science, Networks, Vol 8. Chapter 9. Amsterdam: North Holland.

Sprumont Y. (1990). Population monotonic allocation schemes for cooperative games with transferable utility. Games and Economic Behavior 2: 378-394.

NOTE DI LAVORO DELLA FONDAZIONE ENI ENRICO MATTEI

Fondazione Eni Enrico Mattei Working Paper Series

Our Note di Lavoro are available on the Internet at the following addresses:

<http://www.feem.it/Feem/Pub/Publications/WPapers/default.html>

<http://www.ssrn.com/link/feem.html>

NOTE DI LAVORO PUBLISHED IN 2003

PRIV	1.2003	<i>Gabriella CHIESA and Giovanna NICODANO: <u>Privatization and Financial Market Development: Theoretical Issues</u></i>
PRIV	2.2003	<i>Ibolya SCHINDELE: <u>Theory of Privatization in Eastern Europe: Literature Review</u></i>
PRIV	3.2003	<i>Wietze LISE, Claudia KEMFERT and Richard S.J. TOL: <u>Strategic Action in the Liberalised German Electricity Market</u></i>
CLIM	4.2003	<i>Laura MARSILIANI and Thomas I. RENSTRÖM: <u>Environmental Policy and Capital Movements: The Role of Government Commitment</u></i>
KNOW	5.2003	<i>Reyer GERLAGH: <u>Induced Technological Change under Technological Competition</u></i>
ETA	6.2003	<i>Efrem CASTELNUOVO: <u>Squeezing the Interest Rate Smoothing Weight with a Hybrid Expectations Model</u></i>
SIEV	7.2003	<i>Anna ALBERINI, Alberto LONGO, Stefania TONIN, Francesco TROMBETTA and Margherita TURVANI: <u>The Role of Liability, Regulation and Economic Incentives in Brownfield Remediation and Redevelopment: Evidence from Surveys of Developers</u></i>
NRM	8.2003	<i>Elissaios POPYRAKIS and Reyner GERLAGH: <u>Natural Resources: A Blessing or a Curse?</u></i>
CLIM	9.2003	<i>A. CAPARRÓS, J.-C. PEREAU and T. TAZDAÏT: <u>North-South Climate Change Negotiations: a Sequential Game with Asymmetric Information</u></i>
KNOW	10.2003	<i>Giorgio BRUNELLO and Daniele CHECCHI: <u>School Quality and Family Background in Italy</u></i>
CLIM	11.2003	<i>Efrem CASTELNUOVO and Marzio GALEOTTI: <u>Learning By Doing vs Learning By Researching in a Model of Climate Change Policy Analysis</u></i>
KNOW	12.2003	<i>Carole MAIGNAN, Gianmarco OTTAVIANO and Dino PINELLI (eds.): <u>Economic Growth, Innovation, Cultural Diversity: What are we all talking about? A critical survey of the state-of-the-art</u></i>
KNOW	13.2003	<i>Carole MAIGNAN, Gianmarco OTTAVIANO, Dino PINELLI and Francesco RULLANI (lix): <u>Bio-Ecological Diversity vs. Socio-Economic Diversity. A Comparison of Existing Measures</u></i>
KNOW	14.2003	<i>Maddy JANSSENS and Chris STEYAERT (lix): <u>Theories of Diversity within Organisation Studies: Debates and Future Trajectories</u></i>
KNOW	15.2003	<i>Tuzin BAYCAN LEVENT, Enno MASUREL and Peter NIJKAMP (lix): <u>Diversity in Entrepreneurship: Ethnic and Female Roles in Urban Economic Life</u></i>
KNOW	16.2003	<i>Alexandra BITUSIKOVA (lix): <u>Post-Communist City on its Way from Grey to Colourful: The Case Study from Slovakia</u></i>
KNOW	17.2003	<i>Billy E. VAUGHN and Katarina MLEKOV (lix): <u>A Stage Model of Developing an Inclusive Community</u></i>
KNOW	18.2003	<i>Selma van LONDEN and Arie de RUIJTER (lix): <u>Managing Diversity in a Globalizing World</u></i>
Coalition		
Theory	19.2003	<i>Sergio CURRARINI: <u>On the Stability of Hierarchies in Games with Externalities</u></i>
Network		
PRIV	20.2003	<i>Giacomo CALZOLARI and Alessandro PAVAN (lx): <u>Monopoly with Resale</u></i>
PRIV	21.2003	<i>Claudio MEZZETTI (lx): <u>Auction Design with Interdependent Valuations: The Generalized Revelation Principle, Efficiency, Full Surplus Extraction and Information Acquisition</u></i>
PRIV	22.2003	<i>Marco LiCalzi and Alessandro PAVAN (lx): <u>Tilting the Supply Schedule to Enhance Competition in Uniform-Price Auctions</u></i>
PRIV	23.2003	<i>David ETTINGER (lx): <u>Bidding among Friends and Enemies</u></i>
PRIV	24.2003	<i>Hannu VARTIAINEN (lx): <u>Auction Design without Commitment</u></i>
PRIV	25.2003	<i>Matti KELOHARJU, Kjell G. NYBORG and Kristian RYDQVIST (lx): <u>Strategic Behavior and Underpricing in Uniform Price Auctions: Evidence from Finnish Treasury Auctions</u></i>
PRIV	26.2003	<i>Christine A. PARLOUR and Uday RAJAN (lx): <u>Rationing in IPOs</u></i>
PRIV	27.2003	<i>Kjell G. NYBORG and Ilya A. STREBULAIEV (lx): <u>Multiple Unit Auctions and Short Squeezes</u></i>
PRIV	28.2003	<i>Anders LUNANDER and Jan-Eric NILSSON (lx): <u>Taking the Lab to the Field: Experimental Tests of Alternative Mechanisms to Procure Multiple Contracts</u></i>
PRIV	29.2003	<i>TangaMcDANIEL and Karsten NEUHOFF (lx): <u>Use of Long-term Auctions for Network Investment</u></i>
PRIV	30.2003	<i>Emiel MAASLAND and Sander ONDERSTAL (lx): <u>Auctions with Financial Externalities</u></i>
ETA	31.2003	<i>Michael FINUS and Bianca RUNDSHAGEN: <u>A Non-cooperative Foundation of Core-Stability in Positive Externality NTU-Coalition Games</u></i>
KNOW	32.2003	<i>Michele MORETTO: <u>Competition and Irreversible Investments under Uncertainty</u></i>
PRIV	33.2003	<i>Philippe QUIRION: <u>Relative Quotas: Correct Answer to Uncertainty or Case of Regulatory Capture?</u></i>
KNOW	34.2003	<i>Giuseppe MEDA, Claudio PIGA and Donald SIEGEL: <u>On the Relationship between R&D and Productivity: A Treatment Effect Analysis</u></i>
ETA	35.2003	<i>Alessandra DEL BOCA, Marzio GALEOTTI and Paola ROTA: <u>Non-convexities in the Adjustment of Different Capital Inputs: A Firm-level Investigation</u></i>

GG	36.2003	<i>Matthieu GLACHANT</i> : <u>Voluntary Agreements under Endogenous Legislative Threats</u>
PRIV	37.2003	<i>Narjess BOUBAKRI, Jean-Claude COSSET and Omrane GUEDHAMI</i> : <u>Postprivatization Corporate Governance: the Role of Ownership Structure and Investor Protection</u>
CLIM	38.2003	<i>Rolf GOLOMBEK and Michael HOEL</i> : <u>Climate Policy under Technology Spillovers</u>
KNOW	39.2003	<i>Slim BEN YOUSSEF</i> : <u>Transboundary Pollution, R&D Spillovers and International Trade</u>
CTN	40.2003	<i>Carlo CARRARO and Carmen MARCHIORI</i> : <u>Endogenous Strategic Issue Linkage in International Negotiations</u>
KNOW	41.2003	<i>Sonia OREFFICE</i> : <u>Abortion and Female Power in the Household: Evidence from Labor Supply</u>
KNOW	42.2003	<i>Timo GOESCHL and Timothy SWANSON</i> : <u>On Biology and Technology: The Economics of Managing Biotechnologies</u>
ETA	43.2003	<i>Giorgio Busetti and Matteo MANERA</i> : <u>STAR-GARCH Models for Stock Market Interactions in the Pacific Basin Region, Japan and US</u>
CLIM	44.2003	<i>Katrin MILLOCK and Céline NAUGES</i> : <u>The French Tax on Air Pollution: Some Preliminary Results on its Effectiveness</u>
PRIV	45.2003	<i>Bernardo BORTOLOTTI and Paolo PINOTTI</i> : <u>The Political Economy of Privatization</u>
SIEV	46.2003	<i>Elbert DIJKGRAAF and Herman R.J. VOLLEBERGH</i> : <u>Burn or Bury? A Social Cost Comparison of Final Waste Disposal Methods</u>
ETA	47.2003	<i>Jens HORBACH</i> : <u>Employment and Innovations in the Environmental Sector: Determinants and Econometrical Results for Germany</u>
CLIM	48.2003	<i>Lori SNYDER, Nolan MILLER and Robert STAVINS</i> : <u>The Effects of Environmental Regulation on Technology Diffusion: The Case of Chlorine Manufacturing</u>
CLIM	49.2003	<i>Lori SNYDER, Robert STAVINS and Alexander F. WAGNER</i> : <u>Private Options to Use Public Goods. Exploiting Revealed Preferences to Estimate Environmental Benefits</u>
CTN	50.2003	<i>László Á. KÓCZY and Luc LAUWERS</i> (Ixi): <u>The Minimal Dominant Set is a Non-Empty Core-Extension</u>
CTN	51.2003	<i>Matthew O. JACKSON</i> (Ixi): <u>Allocation Rules for Network Games</u>
CTN	52.2003	<i>Ana MAULEON and Vincent VANNETELBOSCH</i> (Ixi): <u>Farsightedness and Cautiousness in Coalition Formation</u>
CTN	53.2003	<i>Fernando VEGA-REDONDO</i> (Ixi): <u>Building Up Social Capital in a Changing World: a network approach</u>
CTN	54.2003	<i>Matthew HAAG and Roger LAGUNOFF</i> (Ixi): <u>On the Size and Structure of Group Cooperation</u>
CTN	55.2003	<i>Tajji FURUSAWA and Hideo KONISHI</i> (Ixi): <u>Free Trade Networks</u>
CTN	56.2003	<i>Halis Murat YILDIZ</i> (Ixi): <u>National Versus International Mergers and Trade Liberalization</u>
CTN	57.2003	<i>Santiago RUBIO and Alistair ULPH</i> (Ixi): <u>An Infinite-Horizon Model of Dynamic Membership of International Environmental Agreements</u>
KNOW	58.2003	<i>Carole MAIGNAN, Dino PINELLI and Gianmarco I.P. OTTAVIANO</i> : <u>ICT, Clusters and Regional Cohesion: A Summary of Theoretical and Empirical Research</u>
KNOW	59.2003	<i>Giorgio BELLETTINI and Gianmarco I.P. OTTAVIANO</i> : <u>Special Interests and Technological Change</u>
ETA	60.2003	<i>Ronnie SCHÖB</i> : <u>The Double Dividend Hypothesis of Environmental Taxes: A Survey</u>
CLIM	61.2003	<i>Michael FINUS, Ekko van IERLAND and Robert DELLINK</i> : <u>Stability of Climate Coalitions in a Cartel Formation Game</u>
GG	62.2003	<i>Michael FINUS and Bianca RUNDSHAGEN</i> : <u>How the Rules of Coalition Formation Affect Stability of International Environmental Agreements</u>
SIEV	63.2003	<i>Alberto PETRUCCI</i> : <u>Taxing Land Rent in an Open Economy</u>
CLIM	64.2003	<i>Joseph E. ALDY, Scott BARRETT and Robert N. STAVINS</i> : <u>Thirteen Plus One: A Comparison of Global Climate Policy Architectures</u>
SIEV	65.2003	<i>Edi DEFRANCESCO</i> : <u>The Beginning of Organic Fish Farming in Italy</u>
SIEV	66.2003	<i>Klaus CONRAD</i> : <u>Price Competition and Product Differentiation when Consumers Care for the Environment</u>
SIEV	67.2003	<i>Paulo A.L.D. NUNES, Luca ROSSETTO, Arianne DE BLAEIJ</i> : <u>Monetary Value Assessment of Clam Fishing Management Practices in the Venice Lagoon: Results from a Stated Choice Exercise</u>
CLIM	68.2003	<i>ZhongXiang ZHANG</i> : <u>Open Trade with the U.S. Without Compromising Canada's Ability to Comply with its Kyoto Target</u>
KNOW	69.2003	<i>David FRANTZ</i> (Iix): <u>Lorenzo Market between Diversity and Mutation</u>
KNOW	70.2003	<i>Ercole SORI</i> (Iix): <u>Mapping Diversity in Social History</u>
KNOW	71.2003	<i>Ljiljana DERU SIMIC</i> (Ixi): <u>What is Specific about Art/Cultural Projects?</u>
KNOW	72.2003	<i>Natalya V. TARANOVA</i> (Ixi): <u>The Role of the City in Fostering Intergroup Communication in a Multicultural Environment: Saint-Petersburg's Case</u>
KNOW	73.2003	<i>Kristine CRANE</i> (Ixi): <u>The City as an Arena for the Expression of Multiple Identities in the Age of Globalisation and Migration</u>
KNOW	74.2003	<i>Kazuma MATOBA</i> (Ixi): <u>Glocal Dialogue- Transformation through Transcultural Communication</u>
KNOW	75.2003	<i>Catarina REIS OLIVEIRA</i> (Ixi): <u>Immigrants' Entrepreneurial Opportunities: The Case of the Chinese in Portugal</u>
KNOW	76.2003	<i>Sandra WALLMAN</i> (Ixi): <u>The Diversity of Diversity - towards a typology of urban systems</u>
KNOW	77.2003	<i>Richard PEARCE</i> (Ixi): <u>A Biologist's View of Individual Cultural Identity for the Study of Cities</u>
KNOW	78.2003	<i>Vincent MERK</i> (Ixi): <u>Communication Across Cultures: from Cultural Awareness to Reconciliation of the Dilemmas</u>
KNOW	79.2003	<i>Giorgio BELLETTINI, Carlotta BERTI CERONI and Gianmarco I.P. OTTAVIANO</i> : <u>Child Labor and Resistance to Change</u>
ETA	80.2003	<i>Michele MORETTO, Paolo M. PANTEGHINI and Carlo SCARPA</i> : <u>Investment Size and Firm's Value under Profit Sharing Regulation</u>

IEM	81.2003	<i>Alessandro LANZA, Matteo MANERA and Massimo GIOVANNINI: <u>Oil and Product Dynamics in International Petroleum Markets</u></i>
CLIM	82.2003	<i>Y. Hossein FARZIN and Jinhua ZHAO: <u>Pollution Abatement Investment When Firms Lobby Against Environmental Regulation</u></i>
CLIM	83.2003	<i>Giuseppe DI VITA: <u>Is the Discount Rate Relevant in Explaining the Environmental Kuznets Curve?</u></i>
CLIM	84.2003	<i>Reyer GERLAGH and Wietze LISE: <u>Induced Technological Change Under Carbon Taxes</u></i>
NRM	85.2003	<i>Rinaldo BRAU, Alessandro LANZA and Francesco PIGLIARU: <u>How Fast are the Tourism Countries Growing? The cross-country evidence</u></i>
KNOW	86.2003	<i>Elena BELLINI, Gianmarco I.P. OTTAVIANO and Dino PINELLI: <u>The ICT Revolution: opportunities and risks for the Mezzogiorno</u></i>
SIEV	87.2003	<i>Lucas BRETSCGHER and Sjak SMULDERS: <u>Sustainability and Substitution of Exhaustible Natural Resources. How resource prices affect long-term R&D investments</u></i>
CLIM	88.2003	<i>Johan EYCKMANS and Michael FINUS: <u>New Roads to International Environmental Agreements: The Case of Global Warming</u></i>
CLIM	89.2003	<i>Marzio GALEOTTI: <u>Economic Development and Environmental Protection</u></i>
CLIM	90.2003	<i>Marzio GALEOTTI: <u>Environment and Economic Growth: Is Technical Change the Key to Decoupling?</u></i>
CLIM	91.2003	<i>Marzio GALEOTTI and Barbara BUCHNER: <u>Climate Policy and Economic Growth in Developing Countries</u></i>
IEM	92.2003	<i>A. MARKANDYA, A. GOLUB and E. STRUKOVA: <u>The Influence of Climate Change Considerations on Energy Policy: The Case of Russia</u></i>
ETA	93.2003	<i>Andrea BELTRATTI: <u>Socially Responsible Investment in General Equilibrium</u></i>
CTN	94.2003	<i>Parkash CHANDER: <u>The γ-Core and Coalition Formation</u></i>
IEM	95.2003	<i>Matteo MANERA and Angelo MARZULLO: <u>Modelling the Load Curve of Aggregate Electricity Consumption Using Principal Components</u></i>
IEM	96.2003	<i>Alessandro LANZA, Matteo MANERA, Margherita GRASSO and Massimo GIOVANNINI: <u>Long-run Models of Oil Stock Prices</u></i>
CTN	97.2003	<i>Steven J. BRAMS, Michael A. JONES, and D. Marc KILGOUR: <u>Forming Stable Coalitions: The Process Matters</u></i>
KNOW	98.2003	<i>John CROWLEY, Marie-Cecile NAVES (Ixxiii): <u>Anti-Racist Policies in France. From Ideological and Historical Schemes to Socio-Political Realities</u></i>
KNOW	99.2003	<i>Richard THOMPSON FORD (Ixxiii): <u>Cultural Rights and Civic Virtue</u></i>
KNOW	100.2003	<i>Alaknanda PATEL (Ixxiii): <u>Cultural Diversity and Conflict in Multicultural Cities</u></i>
KNOW	101.2003	<i>David MAY (Ixxiii): <u>The Struggle of Becoming Established in a Deprived Inner-City Neighbourhood</u></i>
KNOW	102.2003	<i>Sébastien ARCAND, Danielle JUTEAU, Sirma BILGE, and Francine LEMIRE (Ixxiii) : <u>Municipal Reform on the Island of Montreal: Tensions Between Two Majority Groups in a Multicultural City</u></i>
CLIM	103.2003	<i>Barbara BUCHNER and Carlo CARRARO: <u>China and the Evolution of the Present Climate Regime</u></i>
CLIM	104.2003	<i>Barbara BUCHNER and Carlo CARRARO: <u>Emissions Trading Regimes and Incentives to Participate in International Climate Agreements</u></i>
CLIM	105.2003	<i>Anil MARKANDYA and Dirk T.G. RÜBBELKE: <u>Ancillary Benefits of Climate Policy</u></i>
NRM	106.2003	<i>Anne Sophie CRÉPIN (Ixiv): <u>Management Challenges for Multiple-Species Boreal Forests</u></i>
NRM	107.2003	<i>Anne Sophie CRÉPIN (Ixiv): <u>Threshold Effects in Coral Reef Fisheries</u></i>
SIEV	108.2003	<i>Sara ANIYAR (Ixiv): <u>Estimating the Value of Oil Capital in a Small Open Economy: The Venezuela's Example</u></i>
SIEV	109.2003	<i>Kenneth ARROW, Partha DASGUPTA and Karl-Göran MÄLER (Ixiv): <u>Evaluating Projects and Assessing Sustainable Development in Imperfect Economies</u></i>
NRM	110.2003	<i>Anastasios XEPAPADEAS and Catarina ROSETA-PALMA (Ixiv): <u>Instabilities and Robust Control in Fisheries</u></i>
NRM	111.2003	<i>Charles PERRINGS and Brian WALKER (Ixiv): <u>Conservation and Optimal Use of Rangelands</u></i>
ETA	112.2003	<i>Jack GOODY (Ixiv): <u>Globalisation, Population and Ecology</u></i>
CTN	113.2003	<i>Carlo CARRARO, Carmen MARCHIORI and Sonia OREFFICE: <u>Endogenous Minimum Participation in International Environmental Treaties</u></i>
CTN	114.2003	<i>Guillaume HAERINGER and Myrna WOODERS: <u>Decentralized Job Matching</u></i>
CTN	115.2003	<i>Hideo KONISHI and M. Utku UNVER: <u>Credible Group Stability in Multi-Partner Matching Problems</u></i>
CTN	116.2003	<i>Somdeb LAHIRI: <u>Stable Matchings for the Room-Mates Problem</u></i>
CTN	117.2003	<i>Somdeb LAHIRI: <u>Stable Matchings for a Generalized Marriage Problem</u></i>
CTN	118.2003	<i>Marita LAUKKANEN: <u>Transboundary Fisheries Management under Implementation Uncertainty</u></i>
CTN	119.2003	<i>Edward CARTWRIGHT and Myrna WOODERS: <u>Social Conformity and Bounded Rationality in Arbitrary Games with Incomplete Information: Some First Results</u></i>
CTN	120.2003	<i>Gianluigi VERNASCA: <u>Dynamic Price Competition with Price Adjustment Costs and Product Differentiation</u></i>
CTN	121.2003	<i>Myrna WOODERS, Edward CARTWRIGHT and Reinhard SELTEN: <u>Social Conformity in Games with Many Players</u></i>
CTN	122.2003	<i>Edward CARTWRIGHT and Myrna WOODERS: <u>On Equilibrium in Pure Strategies in Games with Many Players</u></i>
CTN	123.2003	<i>Edward CARTWRIGHT and Myrna WOODERS: <u>Conformity and Bounded Rationality in Games with Many Players</u></i>
	1000	Carlo CARRARO, Alessandro LANZA and Valeria PAPPONETTI: <u>One Thousand Working Papers</u>

NOTE DI LAVORO PUBLISHED IN 2004

IEM	1.2004	<i>Anil MARKANDYA, Suzette PEDROSO and Alexander GOLUB: <u>Empirical Analysis of National Income and So2 Emissions in Selected European Countries</u></i>
ETA	2.2004	<i>Masahisa FUJITA and Shlomo WEBER: <u>Strategic Immigration Policies and Welfare in Heterogeneous Countries</u></i>
PRA	3.2004	<i>Adolfo DI CARLUCCIO, Giovanni FERRI, Cecilia FRALE and Ottavio RICCHI: <u>Do Privatizations Boost Household Shareholding? Evidence from Italy</u></i>
ETA	4.2004	<i>Victor GINSBURGH and Shlomo WEBER: <u>Languages Disenfranchisement in the European Union</u></i>
ETA	5.2004	<i>Romano PIRAS: <u>Growth, Congestion of Public Goods, and Second-Best Optimal Policy</u></i>
CCMP	6.2004	<i>Herman R.J. VOLLEBERGH: <u>Lessons from the Polder: Is Dutch CO2-Taxation Optimal</u></i>
PRA	7.2004	<i>Sandro BRUSCO, Giuseppe LOPOMO and S. VISWANATHAN (lxv): <u>Merger Mechanisms</u></i>
PRA	8.2004	<i>Wolfgang AUSSENEGG, Pegaret PICHLER and Alex STOMPER (lxv): <u>IPO Pricing with Bookbuilding, and a When-Issued Market</u></i>
PRA	9.2004	<i>Pegaret PICHLER and Alex STOMPER (lxv): <u>Primary Market Design: Direct Mechanisms and Markets</u></i>
PRA	10.2004	<i>Florian ENGLMAIER, Pablo GUILLEN, Loreto LLORENTE, Sander ONDERSTAL and Rupert SAUSGRUBER (lxv): <u>The Chopstick Auction: A Study of the Exposure Problem in Multi-Unit Auctions</u></i>
PRA	11.2004	<i>Bjarne BRENDSTRUP and Harry J. PAARSCH (lxv): <u>Nonparametric Identification and Estimation of Multi-Unit, Sequential, Oral, Ascending-Price Auctions With Asymmetric Bidders</u></i>
PRA	12.2004	<i>Ohad KADAN (lxv): <u>Equilibrium in the Two Player, k-Double Auction with Affiliated Private Values</u></i>
PRA	13.2004	<i>Maarten C.W. JANSSEN (lxv): <u>Auctions as Coordination Devices</u></i>
PRA	14.2004	<i>Gadi FIBICH, Arieh GAVIOUS and Aner SELA (lxv): <u>All-Pay Auctions with Weakly Risk-Averse Buyers</u></i>
PRA	15.2004	<i>Orly SADE, Charles SCHNITZLEIN and Jaime F. ZENDER (lxv): <u>Competition and Cooperation in Divisible Good Auctions: An Experimental Examination</u></i>
PRA	16.2004	<i>Marta STRYSZOWSKA (lxv): <u>Late and Multiple Bidding in Competing Second Price Internet Auctions</u></i>
CCMP	17.2004	<i>Slim Ben YOUSSEF: <u>R&D in Cleaner Technology and International Trade</u></i>
NRM	18.2004	<i>Angelo ANTOCI, Simone BORGHESI and Paolo RUSSU (lxvi): <u>Biodiversity and Economic Growth: Stabilization Versus Preservation of the Ecological Dynamics</u></i>
SIEV	19.2004	<i>Anna ALBERINI, Paolo ROSATO, Alberto LONGO and Valentina ZANATTA: <u>Information and Willingness to Pay in a Contingent Valuation Study: The Value of S. Erasmo in the Lagoon of Venice</u></i>
NRM	20.2004	<i>Guido CANDELA and Roberto CELLINI (lxvii): <u>Investment in Tourism Market: A Dynamic Model of Differentiated Oligopoly</u></i>
NRM	21.2004	<i>Jacqueline M. HAMILTON (lxvii): <u>Climate and the Destination Choice of German Tourists</u></i>
NRM	22.2004	<i>Javier Rey-MAQUIEIRA PALMER, Javier LOZANO IBÁÑEZ and Carlos Mario GÓMEZ GÓMEZ (lxvii): <u>Land, Environmental Externalities and Tourism Development</u></i>
NRM	23.2004	<i>Pius ODUNGA and Henk FOLMER (lxvii): <u>Profiling Tourists for Balanced Utilization of Tourism-Based Resources in Kenya</u></i>
NRM	24.2004	<i>Jean-Jacques NOWAK, Mondher SAHLI and Pasquale M. SGRO (lxvii): <u>Tourism, Trade and Domestic Welfare</u></i>
NRM	25.2004	<i>Riaz SHAREEF (lxvii): <u>Country Risk Ratings of Small Island Tourism Economies</u></i>
NRM	26.2004	<i>Juan Luis EUGENIO-MARTÍN, Noelia MARTÍN MORALES and Riccardo SCARPA (lxvii): <u>Tourism and Economic Growth in Latin American Countries: A Panel Data Approach</u></i>
NRM	27.2004	<i>Raúl Hernández MARTÍN (lxvii): <u>Impact of Tourism Consumption on GDP. The Role of Imports</u></i>
CSRM	28.2004	<i>Nicoletta FERRO: <u>Cross-Country Ethical Dilemmas in Business: A Descriptive Framework</u></i>
NRM	29.2004	<i>Marian WEBER (lxvi): <u>Assessing the Effectiveness of Tradable Landuse Rights for Biodiversity Conservation: an Application to Canada's Boreal Mixedwood Forest</u></i>
NRM	30.2004	<i>Trond BJORN DAL, Phoebe KOUNDOURI and Sean PASCOE (lxvi): <u>Output Substitution in Multi-Species Trawl Fisheries: Implications for Quota Setting</u></i>
CCMP	31.2004	<i>Marzio GALEOTTI, Alessandra GORIA, Paolo MOMBRINI and Evi SPANTIDAKI: <u>Weather Impacts on Natural, Social and Economic Systems (WISE) Part I: Sectoral Analysis of Climate Impacts in Italy</u></i>
CCMP	32.2004	<i>Marzio GALEOTTI, Alessandra GORIA, Paolo MOMBRINI and Evi SPANTIDAKI: <u>Weather Impacts on Natural, Social and Economic Systems (WISE) Part II: Individual Perception of Climate Extremes in Italy</u></i>
CTN	33.2004	<i>Wilson PEREZ: <u>Divide and Conquer: Noisy Communication in Networks, Power, and Wealth Distribution</u></i>
KTHC	34.2004	<i>Gianmarco I.P. OTTAVIANO and Giovanni PERI (lxviii): <u>The Economic Value of Cultural Diversity: Evidence from US Cities</u></i>
KTHC	35.2004	<i>Linda CHAIB (lxviii): <u>Immigration and Local Urban Participatory Democracy: A Boston-Paris Comparison</u></i>
KTHC	36.2004	<i>Franca ECKERT COEN and Claudio ROSSI (lxviii): <u>Foreigners, Immigrants, Host Cities: The Policies of Multi-Ethnicity in Rome. Reading Governance in a Local Context</u></i>
KTHC	37.2004	<i>Kristine CRANE (lxviii): <u>Governing Migration: Immigrant Groups' Strategies in Three Italian Cities – Rome, Naples and Bari</u></i>
KTHC	38.2004	<i>Kiflemariam HAMDE (lxviii): <u>Mind in Africa, Body in Europe: The Struggle for Maintaining and Transforming Cultural Identity - A Note from the Experience of Eritrean Immigrants in Stockholm</u></i>
ETA	39.2004	<i>Alberto CAVALIERE: <u>Price Competition with Information Disparities in a Vertically Differentiated Duopoly</u></i>
PRA	40.2004	<i>Andrea BIGANO and Stef PROOST: <u>The Opening of the European Electricity Market and Environmental Policy: Does the Degree of Competition Matter?</u></i>
CCMP	41.2004	<i>Micheal FINUS (lxix): <u>International Cooperation to Resolve International Pollution Problems</u></i>

KTHC	42.2004	<i>Francesco CRESPI</i> : <u>Notes on the Determinants of Innovation: A Multi-Perspective Analysis</u>
CTN	43.2004	<i>Sergio CURRARINI and Marco MARINI</i> : <u>Coalition Formation in Games without Synergies</u>
CTN	44.2004	<i>Marc ESCRHUELA-VILLAR</i> : <u>Cartel Sustainability and Cartel Stability</u>
NRM	45.2004	<i>Sebastian BERVOETS and Nicolas GRAVEL</i> (lxvi): <u>Appraising Diversity with an Ordinal Notion of Similarity: An Axiomatic Approach</u>
NRM	46.2004	<i>Signe ANTHON and Bo JELLESMARK THORSEN</i> (lxvi): <u>Optimal Afforestation Contracts with Asymmetric Information on Private Environmental Benefits</u>
NRM	47.2004	<i>John MBURU</i> (lxvi): <u>Wildlife Conservation and Management in Kenya: Towards a Co-management Approach</u>
NRM	48.2004	<i>Ekin BIROL, Ágnes GYOVAI and Melinda SMALE</i> (lxvi): <u>Using a Choice Experiment to Value Agricultural Biodiversity on Hungarian Small Farms: Agri-Environmental Policies in a Transitional Economy</u>
CCMP	49.2004	<i>Gernot KLEPPER and Sonja PETERSON</i> : <u>The EU Emissions Trading Scheme. Allowance Prices, Trade Flows, Competitiveness Effects</u>
GG	50.2004	<i>Scott BARRETT and Michael HOEL</i> : <u>Optimal Disease Eradication</u>
CTN	51.2004	<i>Dinko DIMITROV, Peter BORM, Ruud HENDRICKX and Shao CHIN SUNG</i> : <u>Simple Priorities and Core Stability in Hedonic Games</u>
SIEV	52.2004	<i>Francesco RICCI</i> : <u>Channels of Transmission of Environmental Policy to Economic Growth: A Survey of the Theory</u>
SIEV	53.2004	<i>Anna ALBERINI, Maureen CROPPER, Alan KRUPNICK and Nathalie B. SIMON</i> : <u>Willingness to Pay for Mortality Risk Reductions: Does Latency Matter?</u>
NRM	54.2004	<i>Ingo BRÄUER and Rainer MARGGRAF</i> (lxvi): <u>Valuation of Ecosystem Services Provided by Biodiversity Conservation: An Integrated Hydrological and Economic Model to Value the Enhanced Nitrogen Retention in Renaturated Streams</u>
NRM	55.2004	<i>Timo GOESCHL and Tun LIN</i> (lxvi): <u>Biodiversity Conservation on Private Lands: Information Problems and Regulatory Choices</u>
NRM	56.2004	<i>Tom DEDEURWAERDERE</i> (lxvi): <u>Bioprospection: From the Economics of Contracts to Reflexive Governance</u>
CCMP	57.2004	<i>Katrin REHDANZ and David MADDISON</i> : <u>The Amenity Value of Climate to German Households</u>
CCMP	58.2004	<i>Koen SMEKENS and Bob VAN DER ZWAAN</i> : <u>Environmental Externalities of Geological Carbon Sequestration Effects on Energy Scenarios</u>
NRM	59.2004	<i>Valentina BOSETTI, Mariaester CASSINELLI and Alessandro LANZA</i> (lxvii): <u>Using Data Envelopment Analysis to Evaluate Environmentally Conscious Tourism Management</u>
NRM	60.2004	<i>Timo GOESCHL and Danilo CAMARGO IGLIORI</i> (lxvi): <u>Property Rights Conservation and Development: An Analysis of Extractive Reserves in the Brazilian Amazon</u>
CCMP	61.2004	<i>Barbara BUCHNER and Carlo CARRARO</i> : <u>Economic and Environmental Effectiveness of a Technology-based Climate Protocol</u>
NRM	62.2004	<i>Elissaios POPYRAKIS and Reyer GERLAGH</i> : <u>Resource-Abundance and Economic Growth in the U.S.</u>
NRM	63.2004	<i>Györgyi BELA, Györgyi PATAKI, Melinda SMALE and Mariann HAJDÚ</i> (lxvi): <u>Conserving Crop Genetic Resources on Smallholder Farms in Hungary: Institutional Analysis</u>
NRM	64.2004	<i>E.C.M. RUIJGROK and E.E.M. NILLESEN</i> (lxvi): <u>The Socio-Economic Value of Natural Riverbanks in the Netherlands</u>
NRM	65.2004	<i>E.C.M. RUIJGROK</i> (lxvi): <u>Reducing Acidification: The Benefits of Increased Nature Quality. Investigating the Possibilities of the Contingent Valuation Method</u>
ETA	66.2004	<i>Giannis VARDAS and Anastasios XEPAPADEAS</i> : <u>Uncertainty Aversion, Robust Control and Asset Holdings</u>
GG	67.2004	<i>Anastasios XEPAPADEAS and Constadina PASSA</i> : <u>Participation in and Compliance with Public Voluntary Environmental Programs: An Evolutionary Approach</u>
GG	68.2004	<i>Michael FINUS</i> : <u>Modesty Pays: Sometimes!</u>
NRM	69.2004	<i>Trond BJØRNDAL and Ana BRASÃO</i> : <u>The Northern Atlantic Bluefin Tuna Fisheries: Management and Policy Implications</u>
CTN	70.2004	<i>Alejandro CAPARRÓS, Abdelhakim HAMMOUDI and Tarik TAZDAÏT</i> : <u>On Coalition Formation with Heterogeneous Agents</u>
IEM	71.2004	<i>Massimo GIOVANNINI, Margherita GRASSO, Alessandro LANZA and Matteo MANERA</i> : <u>Conditional Correlations in the Returns on Oil Companies Stock Prices and Their Determinants</u>
IEM	72.2004	<i>Alessandro LANZA, Matteo MANERA and Michael MCALEER</i> : <u>Modelling Dynamic Conditional Correlations in WTI Oil Forward and Futures Returns</u>
SIEV	73.2004	<i>Margarita GENIUS and Elisabetta STRAZZERA</i> : <u>The Copula Approach to Sample Selection Modelling: An Application to the Recreational Value of Forests</u>
CCMP	74.2004	<i>Rob DELLINK and Ekko van IERLAND</i> : <u>Pollution Abatement in the Netherlands: A Dynamic Applied General Equilibrium Assessment</u>
ETA	75.2004	<i>Rosella LEVAGGI and Michele MORETTO</i> : <u>Investment in Hospital Care Technology under Different Purchasing Rules: A Real Option Approach</u>
CTN	76.2004	<i>Salvador BARBERÀ and Matthew O. JACKSON</i> (lxx): <u>On the Weights of Nations: Assigning Voting Weights in a Heterogeneous Union</u>
CTN	77.2004	<i>Àlex ARENAS, Antonio CABRALES, Albert DÍAZ-GUILERA, Roger GUIMERA and Fernando VEGA-REDONDO</i> (lxx): <u>Optimal Information Transmission in Organizations: Search and Congestion</u>
CTN	78.2004	<i>Francis BLOCH and Armando GOMES</i> (lxx): <u>Contracting with Externalities and Outside Options</u>

CTN	79.2004	<i>Rabah AMIR, Effrosyni DIAMANTOUDI and Licun XUE</i> (lxx): <u>Merger Performance under Uncertain Efficiency Gains</u>
CTN	80.2004	<i>Francis BLOCH and Matthew O. JACKSON</i> (lxx): <u>The Formation of Networks with Transfers among Players</u>
CTN	81.2004	<i>Daniel DIERMEIER, Hülya ERASLAN and Antonio MERLO</i> (lxx): <u>Bicameralism and Government Formation</u>
CTN	82.2004	<i>Rod GARRATT, James E. PARCO, Cheng-ZHONG QIN and Amnon RAPOPORT</i> (lxx): <u>Potential Maximization and Coalition Government Formation</u>
CTN	83.2004	<i>Kfir ELIAZ, Debraj RAY and Ronny RAZIN</i> (lxx): <u>Group Decision-Making in the Shadow of Disagreement</u>
CTN	84.2004	<i>Sanjeev GOYAL, Marco van der LEIJ and José Luis MORAGA-GONZÁLEZ</i> (lxx): <u>Economics: An Emerging Small World?</u>
CTN	85.2004	<i>Edward CARTWRIGHT</i> (lxx): <u>Learning to Play Approximate Nash Equilibria in Games with Many Players</u>
IEM	86.2004	<i>Finn R. FØRSUND and Michael HOEL</i> : <u>Properties of a Non-Competitive Electricity Market Dominated by Hydroelectric Power</u>
KTHC	87.2004	<i>Elissaios POPYRAKIS and Reyer GERLAGH</i> : <u>Natural Resources, Investment and Long-Term Income</u>
CCMP	88.2004	<i>Marzio GALEOTTI and Claudia KEMFERT</i> : <u>Interactions between Climate and Trade Policies: A Survey</u>
IEM	89.2004	<i>A. MARKANDYA, S. PEDROSO and D. STREIMIKIENE</i> : <u>Energy Efficiency in Transition Economies: Is There Convergence Towards the EU Average?</u>
GG	90.2004	<i>Rolf GOLOMBEK and Michael HOEL</i> : <u>Climate Agreements and Technology Policy</u>
PRA	91.2004	<i>Sergei IZMALKOV</i> (lxv): <u>Multi-Unit Open Ascending Price Efficient Auction</u>
KTHC	92.2004	<i>Gianmarco I.P. OTTAVIANO and Giovanni PERI</i> : <u>Cities and Cultures</u>
KTHC	93.2004	<i>Massimo DEL GATTO</i> : <u>Agglomeration, Integration, and Territorial Authority Scale in a System of Trading Cities. Centralisation versus devolution</u>
CCMP	94.2004	<i>Pierre-André JOUVET, Philippe MICHEL and Gilles ROTILLON</i> : <u>Equilibrium with a Market of Permits</u>
CCMP	95.2004	<i>Bob van der ZWAAN and Reyer GERLAGH</i> : <u>Climate Uncertainty and the Necessity to Transform Global Energy Supply</u>
CCMP	96.2004	<i>Francesco BOSELLO, Marco LAZZARIN, Roberto ROSON and Richard S.J. TOL</i> : <u>Economy-Wide Estimates of the Implications of Climate Change: Sea Level Rise</u>
CTN	97.2004	<i>Gustavo BERGANTIÑOS and Juan J. VIDAL-PUGA</i> : <u>Defining Rules in Cost Spanning Tree Problems Through the Canonical Form</u>

- (lix) This paper was presented at the ENGIME Workshop on “Mapping Diversity”, Leuven, May 16-17, 2002
- (lx) This paper was presented at the EuroConference on “Auctions and Market Design: Theory, Evidence and Applications”, organised by the Fondazione Eni Enrico Mattei, Milan, September 26-28, 2002
- (lxi) This paper was presented at the Eighth Meeting of the Coalition Theory Network organised by the GREQAM, Aix-en-Provence, France, January 24-25, 2003
- (lxii) This paper was presented at the ENGIME Workshop on “Communication across Cultures in Multicultural Cities”, The Hague, November 7-8, 2002
- (lxiii) This paper was presented at the ENGIME Workshop on “Social dynamics and conflicts in multicultural cities”, Milan, March 20-21, 2003
- (lxiv) This paper was presented at the International Conference on “Theoretical Topics in Ecological Economics”, organised by the Abdus Salam International Centre for Theoretical Physics - ICTP, the Beijer International Institute of Ecological Economics, and Fondazione Eni Enrico Mattei – FEEM Trieste, February 10-21, 2003
- (lxv) This paper was presented at the EuroConference on “Auctions and Market Design: Theory, Evidence and Applications” organised by Fondazione Eni Enrico Mattei and sponsored by the EU, Milan, September 25-27, 2003
- (lxvi) This paper has been presented at the 4th BioEcon Workshop on “Economic Analysis of Policies for Biodiversity Conservation” organised on behalf of the BIOECON Network by Fondazione Eni Enrico Mattei, Venice International University (VIU) and University College London (UCL), Venice, August 28-29, 2003
- (lxvii) This paper has been presented at the international conference on “Tourism and Sustainable Economic Development – Macro and Micro Economic Issues” jointly organised by CRENoS (Università di Cagliari e Sassari, Italy) and Fondazione Eni Enrico Mattei, and supported by the World Bank, Sardinia, September 19-20, 2003
- (lxviii) This paper was presented at the ENGIME Workshop on “Governance and Policies in Multicultural Cities”, Rome, June 5-6, 2003
- (lxix) This paper was presented at the Fourth EEP Plenary Workshop and EEP Conference “The Future of Climate Policy”, Cagliari, Italy, 27-28 March 2003
- (lxx) This paper was presented at the 9th Coalition Theory Workshop on "Collective Decisions and Institutional Design" organised by the Universitat Autònoma de Barcelona and held in Barcelona, Spain, January 30-31, 2004

2003 SERIES

CLIM	<i>Climate Change Modelling and Policy</i> (Editor: Marzio Galeotti)
GG	<i>Global Governance</i> (Editor: Carlo Carraro)
SIEV	<i>Sustainability Indicators and Environmental Valuation</i> (Editor: Anna Alberini)
NRM	<i>Natural Resources Management</i> (Editor: Carlo Giupponi)
KNOW	<i>Knowledge, Technology, Human Capital</i> (Editor: Gianmarco Ottaviano)
IEM	<i>International Energy Markets</i> (Editor: Anil Markandya)
CSR	<i>Corporate Social Responsibility and Management</i> (Editor: Sabina Ratti)
PRIV	<i>Privatisation, Regulation, Antitrust</i> (Editor: Bernardo Bortolotti)
ETA	<i>Economic Theory and Applications</i> (Editor: Carlo Carraro)
CTN	<i>Coalition Theory Network</i>

2004 SERIES

CCMP	<i>Climate Change Modelling and Policy</i> (Editor: Marzio Galeotti)
GG	<i>Global Governance</i> (Editor: Carlo Carraro)
SIEV	<i>Sustainability Indicators and Environmental Valuation</i> (Editor: Anna Alberini)
NRM	<i>Natural Resources Management</i> (Editor: Carlo Giupponi)
KTHC	<i>Knowledge, Technology, Human Capital</i> (Editor: Gianmarco Ottaviano)
IEM	<i>International Energy Markets</i> (Editor: Anil Markandya)
CSR	<i>Corporate Social Responsibility and Management</i> (Editor: Sabina Ratti)
PRA	<i>Privatisation, Regulation, Antitrust</i> (Editor: Bernardo Bortolotti)
ETA	<i>Economic Theory and Applications</i> (Editor: Carlo Carraro)
CTN	<i>Coalition Theory Network</i>